

2.20	Estimating the entanglement of proposed class of states using the Bell nonlocality measure M^W , nonlinear entanglement witness E^W and the rescaled FEF F^W as a function of weak measurement strength η , considering $\gamma = 0.6$.	46
2.21	Estimation of fully-entangled fraction for a given nonlocality for proposed class of states, pure states and randomly generated two-qubit states (green area) and the coordinates of points ϱ^k (where $k=1,\dots,6$) to characterize the proposed class of states, given in Table 2.1.	47
2.22	Channel capacity of the proposed class of states ϱ for superdense coding protocol as a function of weak measurement strength η at three different values of decoherence parameter γ .	48
2.23	Comparison of channel capacity of the proposed class with other bipartite entangled mixed states.	48
3.1	A scenario to analyse the effect of decoherence on maximum expectation value of the Svetlichny operator.	54
3.2	Estimation of maximum value of $S_v(\rho_{gA}^\gamma)_{opt}$ vs 3-tangle (τ) of the initial 3-qubit GHZ state with respect to decoherence parameter γ .	56
3.3	Plot of $S_v(\rho_{gA}^\gamma)_{opt}$ with respect to 3-tangle (τ) of the initial 3-qubit GHZ state for four different values of decoherence parameter.	57
3.4	A scenario to analyse the effect of weak measurement and its reversal operations on the existence of genuine tripartite nonlocal correlations.	57
3.5	Estimation of maximum value of $S_v(\rho_{gA}^{wk})_{opt}$ vs 3-tangle (τ) of the initial 3-qubit GHZ state with respect to weak measurement strength parameter η , considering $\gamma = 0.5$.	58
3.6	$S_v(\rho_{gA}^{wk})_{opt}$ with respect to 3-tangle (τ) of initial state for different weak measurement strengths, considering $\gamma = 0.5$.	59
3.7	Effect of weak measurement on the maximum expectation value of the Svetlichny operator as a function of noise parameter γ .	59
3.8	Estimation of maximum value of $S_v(\rho_{gP}^\gamma)_{opt}$ vs 3-tangle (τ) of the initial 3-qubit GHZ state with respect to decoherence parameter γ .	61
3.9	Plot of $S_v(\rho_{gP}^\gamma)_{opt}$ with respect to 3-tangle (τ) of the initial 3-qubit GHZ state for four different values of decoherence parameter.	62
3.10	Plot of $S_v(\rho_{gP}^\gamma)_{opt}$ with respect to decoherence parameter γ for three different input states.	62
3.11	Effect of decoherence on $S_v(\rho_{gP}^{wk})_{opt}$ for a maximally and a partially entangled input state under applications of weak measurement.	64
3.12	Effect of weak measurement on $S_v(\rho_{gP}^{wk})_{opt}$ for a maximally entangled state, considering $\eta_r = 0.6$ at different values of γ .	65
3.13	Effect of weak measurement on $S_v(\rho_{gP}^{wk})_{opt}$ for a initial state with $\theta = \pi/6$, considering $\eta_r = 0.8$ at different values of γ .	65
3.14	Plot of $S_v(\rho_{gD}^\gamma)_{opt}$ with respect to 3-tangle (τ) of the initial 3-qubit GHZ state for three different values of decoherence parameter.	67
3.15	Effect of depolarizing channel on $S_v(\rho_{gD}^\gamma)_{opt}$ for three different initial input states.	67
3.16	Plot of $S_v(\rho_{gD}^{wk})_{opt}$ with respect to 3-tangle (τ) of the initial three-qubit GHZ state for three different values of weak measurement strength, considering $\gamma = 0.1$.	69
3.17	Effect of weak measurement on $S_v(\rho_{gD}^{wk})_{opt}$ for three different initial input states, considering $\gamma = 0.1$.	69
3.18	$S_v(\rho_{gs}^{wk})_{opt}$ with respect to 3-tangle (τ) of the initial state for different weak measurement strengths, considering $\gamma = 0.5$ for two different values of $\theta_3 = (\frac{\pi}{4}, \frac{\pi}{2})$.	71
3.19	$S_v(\rho_{ms}^{wk})_{opt}$ with respect to 3-tangle (τ) of the initial state for different weak measurement strengths, considering $\gamma = 0.5$ and $\theta = \frac{\pi}{4}$.	72
3.20	Negativity of the finally shared 3-qubit state as a function of noise parameter γ .	72
3.21	Negativity of the finally shared 3-qubit state as a function of weak measurement strength η at different values of decoherence parameter γ .	73

3.22	Plot of maximum expectation value of the Svetlichny operator $S_v(\rho_{gs}^{wk})_{opt}$ with respect to Negativity N of the finally shared state for different values of weak measurement strength, considering $\gamma = 0.5$.	73
3.23	Maximum expectation value of $S_v(\rho_W^{wk})_{opt}$ with respect to the varying sum of concurrences $(C_{12} + C_{23} + C_{31})$, for different weak measurement strengths, considering $C_{12} = \frac{2}{3}$ and $\gamma = 0.1$.	75
3.24	$S_v(\rho_{W_n}^{wk})_{opt}$ with respect to the varying sum of concurrences $(C''_{12} + C''_{23} + C''_{31})$, for different weak measurement strengths, considering $\gamma = 0.1$.	76
3.25	Maximum expectation value of $S_v(\rho_{W_n}^{wk})_{opt}$ vs n for W_n type states at different weak measurement strengths, considering $\gamma = 0.1$.	76
3.26	Estimation of $S'_v(\rho_G^\gamma)_{opt}$ with respect to 4-qubit entanglement (τ_4) of the initial GHZ state considering four different values of decoherence parameter γ .	81
3.27	Estimation of $S'_v(\rho_G^{wk})_{opt}$ with respect to 4-qubit entanglement (τ_4) of the initial GHZ state for different weak measurement strengths, considering $\gamma = 0.5$.	81
3.28	Maximum expectation value of the Svetlichny operator as a function of noise parameter γ for the finally shared state at different values of weak measurement strength.	82
3.29	Negativity of the finally shared 4-qubit state as a function of noise parameter γ .	83
3.30	Negativity of the finally shared 4-qubit state as a function of weak measurement strength at three different values of noise parameter γ .	83
3.31	Estimation of $S'_v(\rho_G^{wk})_{opt}$ with respect to Negativity $N(\rho_G^{wk})$ for different values of weak measurement strengths, considering $\gamma = 0.5$.	83
4.1	Estimation of $B_C(\Psi\rangle)_{opt}$ with respect to the state parameter α .	89
4.2	Estimation of $B_C(\varrho)_{opt}$ with respect to state parameter η of the proposed state in chapter-2 and decoherence parameter γ .	92
4.3	A comparison between estimated values of geometric discord, and maximum and minimum value of geometric discord as evaluated using correlation coefficients with respect to the state parameter θ .	95
4.4	A comparison between estimated values of geometric discord, and maximum and minimum value of geometric discord as evaluated using correlation coefficients with respect to the state parameter of the Werner state.	96
4.5	A comparison between estimated values of geometric discord, and maximum and minimum value of geometric discord as evaluated using correlation coefficients with respect to the state parameter of Horodecki states.	96
4.6	A comparison between estimated values of geometric discord, and maximum and minimum value of geometric discord as evaluated using correlation coefficients with respect to the state parameter of the state proposed by WenChao Ma <i>et al.</i>	97
4.7	A comparison between estimated values of geometric discord, and maximum and minimum value of geometric discord as evaluated using correlation coefficients with respect to the state parameter of the proposed class of states in chapter-2.	98
4.8	Estimation of maximum expectation value of $S_C(\rho_g)_{opt}$ with respect to 3-tangle $\tau(\rho_g)$ of three-qubit generalized GHZ states.	100
5.1	Comparison of classical and quantum bounds of CHSH operator varying with parameters p, q in a biased experimental set-up.	107
5.2	Region in $[p, q]$ space where fine-grained uncertainty can distinguish between quantum and classical correlations.	109
5.3	Maximum possible classical and quantum scores of a CHSH game with an arbitrary pure bipartite entangled state for different values of biasing parameters p and q .	112
6.1	Comparison of efficacies of shared bipartite states in three optimal cases.	120
6.2	Comparison of the efficiencies of three and four-qubit W-type states as resources.	122

