

List of Figures

Figures	Title	page
2.1	General representation of ubiquitin proteasome pathway and its enzymes	8
2.2	An illustration of orchestration of factors involved in neurodevelopment and neurodegeneration	10
2.3	Overview of current understanding of amyloids	15
2.4	An overview of orchestration of cellular proteolytic machinery to reestablish homeostasis	19
3.1	<i>Lanosterol treatment upregulates cellular autophagy and induces co-chaperone CHIP expression</i>	22
3.2	<i>Treatment of lanosterol enhances the mRNA expression level of CHIP</i>	23
3.3	<i>Time-dependent treatment of lanosterol increases cellular level of CHIP protein</i>	24
3.4	<i>Lanosterol modulates CHIP levels in concentration-dependent manner</i>	24
3.5	<i>Expression of GFP-wtCAT plasmid constructs</i>	25
3.6	<i>Effect of chloroquine treatment on the expression of GFP-wtCAT</i>	25
3.7	<i>Lanosterol enhances the clearance of a mutant bona-fide misfolded proteins</i>	26
3.8	<i>Chloroquine treatment enhances the aggregation propensity of modeled misfolded protein</i>	26
3.9	<i>Lanosterol reduces the intracellular perinuclear accumulation of heat-denatured luciferase protein</i>	27
3.10	<i>Effect of lanosterol on the degradation of heat-denatured luciferase protein</i>	28
3.11	<i>Effects of lanosterol treatment on overall cellular health</i>	29
3.12	<i>Expression of SOD1-WT plasmid constructs</i>	30
3.13	<i>Effects of autophagy inhibition on the aggregation of ubiquitinated cellular proteins</i>	30
3.14	<i>Lanosterol suppresses the mutant SOD1 aggresomes formation in cells</i>	31
3.15	<i>Higher accumulation of mutant SOD1-G37R protein following autophagy inhibition</i>	31
3.16	<i>Wild-type alpha-synuclein expression analysis</i>	32
3.17	<i>Effects of administration of chloroquine on the expression of α-synuclein</i>	32
3.18	<i>Treatment of lanosterol reduces aggregation of mutant α-synuclein</i>	33
3.19	<i>Inhibition of autophagy overturns the anti-aggregatory effects of lanosterol</i>	33
3.20	<i>Representation of interaction of lanosterol with CHIP TPR domain</i>	34
3.21	<i>Lanosterol stabilizes co-chaperone CHIP</i>	35
3.22	<i>Expression of ataxin-3(Q28) plasmid constructs</i>	35
3.23	<i>Inhibition of autophagy leads to increased accumulation of normal ataxin-3(Q28) containing proteins</i>	36
3.24	<i>Lanosterol induces suppression of pathogenic ataxin-3 polyglutamine protein aggregation in cells</i>	36
3.25	<i>Administration of chloroquine ameliorates the cytoprotective effects of lanosterol</i>	37
3.26	<i>Visualization of normal huntingtin protein expression in Cos-7 cells</i>	38
3.27	<i>Effects of inhibition of cellular autophagic flux on the expression of normal huntingtin protein</i>	38
3.28	<i>Treatment of lanosterol results in decrease of cytoplasmic misfolded inclusions of expanded huntingtin polyglutamine aggregates</i>	39
3.29	<i>Monitoring of impact of chloroquine treatment on the formation of perinuclear inclusions of polyglutamine-containing huntingtin inside the cells</i>	39
3.30	<i>Lanosterol reduces misfolded protein aggregation</i>	40
3.31	<i>Lanosterol alleviates stress-induced cytotoxicity</i>	41
3.32	<i>Suppression of overall cytoplasmic aggregate formation following lanosterol treatment</i>	42
4.1	<i>Ibuprofen treatment causes accumulation of ubiquitinated proteins</i>	48
4.2	<i>Visualization of morphological changes caused by ibuprofen treatment</i>	48
4.3	<i>Treatment of ibuprofen disturbs proteasome function and induces cytotoxicity</i>	49
4.4	<i>Exposure of ibuprofen induces loss of chymotrypsin-like proteasome activity in dose-dependent manner</i>	50
4.5	<i>Time-dependent loss of chymotrypsin-like proteasomal activity in response to ibuprofen administration</i>	50
4.6	<i>Ibuprofen also suppresses post-glutamyl peptide hydrolase-like activity of proteasome</i>	51

4.7	<i>Time-dependent inhibition of post-glutamyl peptide hydrolase activity of 20S proteasome</i>	51
4.8	<i>Ibuprofen treatment leads to stabilization of misfolded proteins in cells</i>	52
4.9	<i>Effect of ibuprofen treatment on soluble fraction of pathogenic huntingtin protein</i>	53
4.10	<i>Ibuprofen treatment stabilizes ubiquitinated proteins inside the cells</i>	54
4.11	<i>Concentration-dependent cytotoxicity conferred by ibuprofen</i>	54
4.12	<i>Ibuprofen presents in silico interaction with proteasomal subunits</i>	55
4.13	<i>Enhanced accumulation of cytoplasmic ubiquitinated proteins following ibuprofen-mediated proteasomal inhibition</i>	56
4.14	<i>Bright-field image micrographs of ibuprofen treated cells</i>	56
4.15	<i>Formation of aggresomes of misfolded proteins response to ibuprofen treatment in cells</i>	57
4.16	<i>Ibuprofen mediates stabilization of IκB-α</i>	58
4.17	<i>Downregulation of NF-κB transcriptional activity in response to ibuprofen treatment</i>	58
4.18	<i>Ibuprofen causes stabilization of aggregation-prone misfolded protein species</i>	59
4.19	<i>Treatment of ibuprofen accelerates the accumulation of proteasome and ubiquitin-positive protein aggregates</i>	59
4.20	<i>Proteasomal inhibition by ibuprofen leads to increased accumulation of cell cycle regulatory proteins p53 and cyclin-dependent kinase inhibitor p27</i>	60
4.21	<i>Ibuprofen induces cytoplasmic accumulation of pro-apoptotic proteasomal target proteins</i>	60
4.22	<i>Ibuprofen treatment induces apoptosis and causes nuclear morphological changes</i>	61
4.23	<i>Time-dependent assessment of apoptosis by flow cytometry using annexin V-FITC and propidium iodide double staining</i>	62
4.24	<i>Representation of DNA fragmentation following treatment of inbuprofen</i>	63
4.25	<i>Ibuprofen treatment induces cytochrome c release during apoptosis</i>	64
4.26	<i>Higher accumulation of cytochrome c following the treatment of cells with ibuprofen</i>	64
4.27	<i>Ibuprofen treatment leads to mitochondrial depolarization</i>	65
4.28	<i>Analysis of ibuprofen-mediated mitochondrial alterations using JC-1 staining and fluorescence analysis</i>	65