

## List of Figures

Figures	Title	Page
1.1	<i>Innate immune sensing is mediated via germline-encoded pattern recognition receptors</i>	1
2.1	<i>Crosstalk between cancer cells and innate immune cells forms the highly enriched tumor microenvironment.</i>	4
2.2	<i>Inflammasome activation pathways.</i>	5
2.3	<i>Current therapeutic advances present NLRs as promising drug targets for fostering host innate immune responses against cancer.</i>	23
3.1	<i>Amorphous nanosilica induces ASC-mediated inflammation in human lung cell population.</i>	26
3.2	<i>Characterization of amorphous silica nanoparticles by transmission electron microscopy.</i>	29
3.3	<i>Schematic for experimental design</i>	30
3.4	<i>MTT Assay for assessment of cell viability following nanosilica exposure.</i>	31
3.5	<i>Amorphous nanosilica dependent cytotoxicity is Caspase-1 and NLRP3 independent.</i>	32
3.6	<i>ASC expression in nanosilica treated cells was analysed by immunofluorescence.</i>	33
3.7	<i>Confocal Microscopy for ASC speck quantification in epithelial cells</i>	34
3.8	<i>Characterization of nanosilica internalization and subsequent cell death</i>	35
3.9	<i>Cytokine expression analysis in response to nanosilica exposure</i>	36
4.1	<i>Schematic flow diagram of multidimensional investigation exploring the role of specialized nucleotide-binding domain and leucine rich-repeat containing receptors (NLRs) in glioma pathology.</i>	39
4.2	<i>NLR gene expression pattern in glioblastoma using TCGA database.</i>	43
4.3	<i>NLR gene expression pattern in low grade glioma using TCGA database.</i>	44
4.4	<i>NLR gene expression networks in low grade glioma and glioblastoma.</i>	45
4.5	<i>Heat map clustering for NLR gene expression in glioma.</i>	47
4.6	<i>Negative regulation of NLR gene expression by methylation in glioblastoma.</i>	52
4.7	<i>Gene ontology (GO) analysis of significantly altered genes and associated biological processes.</i>	53
4.8	<i>Gene set enrichment analysis (GSEA) delineates gene ontology (GO) of significantly altered genes and their interactions within the biological reactome.</i>	54
4.9	<i>Kaplan-Meier (KM) survival curves of TCGA low grade glioma (grade 2) stratified by the expression of NLRs.</i>	56
4.10	<i>Kaplan-Meier (KM) survival curves of TCGA low grade glioma (grade 3) and glioblastoma (grade 4) patients stratified by the expression levels of NLRs.</i>	57
4.11	<i>Kaplan-Meier (KM) survival curves for glioblastoma and low grade glioma stratified by the expression of NLRs and NLR-associated genes using REMBRANDT.</i>	59
4.12	<i>Immunocytochemistry for ASC protein expression in microglia and macrophages.</i>	61
4.13	<i>Immunocytochemistry for ASC protein expression in gliomas.</i>	62
4.14	<i>Immunocytochemistry for AIM2 protein expression in microglia and macrophages.</i>	63
4.15	<i>Immunocytochemistry for AIM2 protein expression in gliomas.</i>	64
4.16	<i>Immunocytochemistry for ASC and AIM2 protein expression in primary endothelial cells.</i>	65
4.17	<i>Immunocytochemistry for Caspase-1 protein expression in glioma and microglial cells.</i>	66
4.18	<i>Immunocytochemistry for NLRP-12 protein expression in glioma and microglial cells.</i>	67
4.19	<i>NLRP12 inhibition increases cellular proliferation in vitro</i>	68
5.1	<i>Analytic workflow for study of fenugreek-induced anti-inflammatory effects and possible interactions with ASC protein targeting in vitro and in silico approaches.</i>	71
5.2	<i>Fenugreek increases ASC speck formation in nanosilica treated cells.</i>	72
5.3	<i>Fenugreek seed extract inhibits nanosilica-induced cytotoxicity.</i>	73