

## List of Tables

<i>Table</i>	<i>Title</i>	<i>Page</i>
2.1	Masses and charges of atoms of BTMAC and SA used in the atomistic simulations. These parameters are taken from the previously studied system [Debnath <i>et al.</i> , 2009].	21
2.2	Number of molecules of BTMAC, SA and water used in different water concentrations investigated. BTMAC to SA ratio is fixed at 2 : 1. The percentage of water decreases from S1 to S2.	21
2.3	Number of molecules of BTMAC, SA and water in different systems investigated. BTMAC to SA ratio is fixed at 2 : 1 in A1 and A2. The percentage of water decreases from A1 to A2.	22
3.1	Number of molecules in systems A1, B1 and B11.	35
3.2	Definition of bead type in the MARTINI model	36
3.3	Mapping scheme for coarse-grained simulation	36
3.4	Non-bonded parameters used in CG B11 where $M$ , $M_a$ and $M_b$ represent the MARTINI model and their modifications respectively. $PM$ is the polarizable MARTINI water model.	39
4.1	Compositions, initial and final configurations, run-lengths and asymmetry percentage of systems investigated. The asymmetry in each system is computed using equation 4.1. The asymmetry percentage remains unchanged from AA to CG scales except S5 which has a standard deviation of 0.77% in asymmetry.	48
4.2	Area per surfactant molecule (APS) ( $nm^2$ ) and volume per surfactant (VPS) ( $nm^3$ ) of simulated bilayers at AA and CG resolutions. $M_p$ and $L_p$ denoted the more and the less populated leaflets respectively. Standard deviations of APS at AA and CG resolutions are 0.001 and 0.0002-0.0003 respectively. <sup>a</sup> [Lunkad <i>et al.</i> , 2017], <sup>b</sup> [N Stelter and Keyes, 2017], <sup>c</sup> [Boek <i>et al.</i> , 2005a].	49
4.3	Extent of interdigitation ( $O_{overlap}$ ) of simulated bilayers at AA and CG resolutions. S6 is prepared only for CG runs as the system size is large.	53
4.4	Bilayer thickness (in nm) from AA and CG simulations. <sup>a</sup> [Lunkad <i>et al.</i> , 2017], <sup>b</sup> [Ritwiset <i>et al.</i> , 2016], <sup>c</sup> [Khakbaz and Klauda, 2018], <sup>d</sup> [Zhao <i>et al.</i> , 2019], <sup>e</sup> [Takahashi <i>et al.</i> , 1997].	60
4.5	Average single chain configurational entropy, $S_{conf}^{per-chain}$ , ( $JK^1mol^{-1}$ ) with standard deviations calculated from time-frames over which $S_{conf}^{per-chain}$ converges and all molecules of BTMAC and SA in AA and CG simulations. All the values are rounded off to their nearest integers and the standard deviations smaller than 0.1 are not reported here.	62
4.6	Average order parameter, $ S_{CD} $ of simulated bilayers at AA resolutions. <sup>a</sup> [Debnath <i>et al.</i> , 2009], <sup>b</sup> [Lunkad <i>et al.</i> , 2017], <sup>c</sup> [Khakbaz and Klauda, 2018], <sup>d</sup> [Zhao <i>et al.</i> , 2019], <sup>e</sup> [Davis, 1979]	70
4.7	Average order parameter, $P_2$ of simulated bilayers at AA and CG resolutions.	71
5.1	Binding energies ( $kcal\ mol^{-1}$ ), angles ( $^\circ$ ) and distances ( $\text{\AA}$ ) for the RH and LH dimers. The results are compared with reported values for similar PDI analogues and PDI derivatives. [Ahmed <i>et al.</i> , 2017] <sup>a</sup> , [Oltean <i>et al.</i> , 2013] <sup>b</sup> , [Chen <i>et al.</i> , 2015] <sup>c</sup> .	84
5.2	Non-bonded energies, closest distance and inter-planar angle of dimers with positive (RH) and negative (LH) inter- planar angle from molecular dynamics simulations.	88

6.1	Energies of the open and folded dimers, stacked dimers and their stacking energies obtained from the DFTB calculations. The folded PyKC conformer has a stronger stacking energy. The inter-planar distances obtained after DFTB geometry optimization and PXRD experiment [Singha <i>et al.</i> , 2019] are in good agreement indicating towards the presence of $\pi - \pi$ stacking.	100
A1	Box lengths of the simulated bilayers for AA and CG simulations.	107