

References

- Agarwal, S. and Sohal, R.S. (1994), "Aging and protein oxidative damage", *Mechanisms of Ageing and Development*, Vol. 75, No. 1, pp. 11-19
- Aguzzi, A. and Steele, A.D. (2009), "Prion topology and toxicity", *Cell*, Vol. 137, No. 6, pp. 994-996
- Amati, B. (2004), "Myc degradation: Dancing with Ubiquitin ligases", *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 101, No. 24, pp. 8843-8844
- Apodaca, J., Kim, I. and Rao, H. (2006), "Cellular tolerance of prion protein PrP in yeast involves proteolysis and the unfolded protein response", *Biochemical and Biophysical Research Communications*, Vol. 347, No. 1, pp. 319-326
- Ardley, H.C., Scott, G.B., Rose, S.A., Tan, N.G.S., Markham, A.F. and Robinson, P.A. (2003), "Inhibition of Proteasomal Activity Causes Inclusion Formation in Neuronal and Non-Neuronal Cells Overexpressing Parkin", *Molecular Biology of the Cell*, Vol. 14, No. 11, pp. 4541-4556
- Ballinger, C.A., Connell, P., Wu, Y., Hu, Z., Thompson, L.J., Yin, L.Y. and Patterson, C. (1999), "Identification of CHIP, a novel tetratricopeptide repeat-containing protein that interacts with heat shock proteins and negatively regulates chaperone functions", *Molecular and Cellular Biology*, Vol. 19, No. 6, pp. 4535-4545
- Bandopadhyay, R., Kingsbury, A.E., Muqit, M.M., Harvey, K., Reid, A.R., Kilford, L., Engelender, S., Schlossmacher, M.G., Wood, N.W., Latchman, D.S., Harvey, R.J. and Lees, A.J. (2005), "Synphilin-1 and parkin show overlapping expression patterns in human brain and form aggresomes in response to proteasomal inhibition", *Neurobiology of Disease*, Vol. 20, No. 2, pp. 401-411
- Bashan, A. and Yonath, A. (2008), "Correlating ribosome function with high-resolution structures", *Trends in Microbiology*, Vol. 16, No. 7, pp. 326-335
- Bazan, N.G., Eady, T.N., Khoutorova, L., Atkins, K.D., Hong, S., Lu, Y., Zhang, C., Jun, B., Obenaus, A., Fredman, G., Zhu, M., Winkler, J.W., Petasis, N.A., Serhan, C.N. and Belayev, L. (2012), "Novel aspirin-triggered neuroprotectin D1 attenuates cerebral ischemic injury after experimental stroke", *Experimental Neurology*, Vol. 236, No. 1, pp. 122-130
- Beal, M.F. (2005), "Mitochondria take center stage in aging and neurodegeneration", *Annals of Neurology*, Vol. 58, No. 4, pp. 495-505
- Bejarano, E. and Cuervo, A.M. (2010), "Chaperone-mediated autophagy", *Proceedings of the American Thoracic Society*, Vol. 7, No. 1, pp. 29-39
- Bence, N.F., Sampat, R.M. and Kopito, R.R. (2001), "Impairment of the Ubiquitin-proteasome system by protein aggregation", *Science*, Vol. 292, No. 5521, pp. 1552-1555
- Bengtson, M.H. and Joazeiro, C.A.P. (2010), "Role of a ribosome-associated E3 Ubiquitin ligase in protein quality control", *Nature*, Vol. 467, No. 7314, pp. 470-473
- Bennett, E.J., Bence, N.F., Jayakumar, R. and Kopito, R.R. (2005), "Global impairment of the Ubiquitin-proteasome system by nuclear or cytoplasmic protein aggregates precedes inclusion body formation", *Molecular Cell*, Vol. 17, No. 3, pp. 351-365
- Bercovich, B., Stancovski, I., Mayer, A., Blumenfeld, N., Laszlo, A., Schwartz, A.L. and Ciechanover, A. (1997), "Ubiquitin-dependent degradation of certain protein substrates in vitro requires the molecular chaperone Hsc70", *Journal of Biological Chemistry*, Vol. 272, No. 14, pp. 9002-9010
- Berke, S.J.S. and Paulson, H.L. (2003), "Protein aggregation and the Ubiquitin proteasome pathway: Gaining the UPPER hand on neurodegeneration", *Current Opinion in Genetics and Development*, Vol. 13, No. 3, pp. 253-261
- Bett, J.S., Goellner, G.M., Woodman, B., Pratt, G., Rechsteiner, M. and Bates, G.P. (2006), "Proteasome impairment does not contribute to pathogenesis in R6/2 Huntington's disease mice: exclusion of proteasome activator REGgamma as a therapeutic target", *Human Molecular Genetics*, Vol. 15, No. 1, pp. 33-44

- Bonifati, V., Rizzu, P., Van Baren, M.J., Schaap, O., Breedveld, G.J., Krieger, E., Dekker, M.C.J., Squitieri, F., Ibanez, P., Joosse, M., Van Dongen, J.W., Vanacore, N., Van Swieten, J.C., Brice, A., Meco, G., Van Duijn, C.M., Oostra, B.A. and Heutink, P. (2003), "Mutations in the DJ-1 gene associated with autosomal recessive early-onset parkinsonism", *Science*, Vol. 299, No. 5604, pp. 256-259
- Bradford, M.M. (1976), "A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding", *Analytical Biochemistry*, Vol. 72, No., pp. 248-254
- Brandt, F., Etchells, S.A., Ortiz, J.O., Elcock, A.H., Hartl, F.U. and Baumeister, W. (2009), "The Native 3D Organization of Bacterial Polysomes", *Cell*, Vol. 136, No. 2, pp. 261-271
- Breckenridge, D.G., Germain, M., Mathai, J.P., Nguyen, M. and Shore, G.C. (2003), "Regulation of apoptosis by endoplasmic reticulum pathways", *Oncogene*, Vol. 22, No. 53 REV. ISS. 7, pp. 8608-8618
- Burnett, B., Li, F. and Pittman, R.N. (2003), "The polyglutamine neurodegenerative protein ataxin-3 binds polyubiquitylated proteins and has Ubiquitin protease activity", *Human Molecular Genetics*, Vol. 12, No. 23, pp. 3195-3205
- Burnett, B.G. and Pittman, R.N. (2005), "The polyglutamine neurodegenerative protein ataxin 3 regulates aggresome formation", *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 102, No. 12, pp. 4330-4335
- Burns, M.P., Zhang, L., Rebeck, G.W., Querfurth, H.W. and Moussa, C.E.H. (2009), "Parkin promotes intracellular A β 1-42 clearance", *Human Molecular Genetics*, Vol. 18, No. 17, pp. 3206-3216
- Burr, M.L., Cano, F., Svobodova, S., Boyle, L.H., Boname, J.M. and Lehner, P.J. (2011), "HRD1 and UBE2J1 target misfolded MHC class I heavy chains for endoplasmic reticulum-associated degradation", *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 108, No. 5, pp. 2034-2039
- Calandria, J.M., Mukherjee, P.K., De Rivero Vaccari, J.C., Zhu, M., Petasis, N.A. and Bazan, N.G. (2012), "Ataxin-1 poly(Q)-induced proteotoxic stress and apoptosis are attenuated in neural cells by docosahexaenoic acid-derived neuroprotectin D1", *Journal of Biological Chemistry*, Vol. 287, No. 28, pp. 23726-23739
- Callow, M.G., Tran, H., Phu, L., Lau, T., Lee, J., Sandoval, W.N., Liu, P.S., Bheddah, S., Tao, J., Lill, J.R., Hongo, J.A., Davis, D., Kirkpatrick, D.S., Polakis, P. and Costa, M. (2011), "Ubiquitin ligase RNF146 regulates tankyrase and Axin to promote Wnt signaling", *PLoS ONE*, Vol. 6, No. 7,
- Carrell, R.W. and Lomas, D.A. (1997), "Conformational disease", *Lancet*, Vol. 350, No. 9071, pp. 134-138
- Carvalho, P., Goder, V. and Rapoport, T.A. (2006), "Distinct Ubiquitin-Ligase Complexes Define Convergent Pathways for the Degradation of ER Proteins", *Cell*, Vol. 126, No. 2, pp. 361-373
- Chakrabarti, O. and Hegde, R.S. (2009), "Functional Depletion of Mahogunin by Cytosolically Exposed Prion Protein Contributes to Neurodegeneration", *Cell*, Vol. 137, No. 6, pp. 1136-1147
- Chan, E.M., Young, E.J., Ianzano, L., Munteanu, I., Zhao, X., Christopoulos, C.C., Avanzini, G., Elia, M., Ackerley, C.A., Jovic, N.J., Bohlega, S., Andermann, E., Rouleau, G.A., Delgado-Escueta, A.V., Minassian, B.A. and Scherer, S.W. (2003), "Mutations in NHLRC1 cause progressive myoclonus epilepsy", *Nature Genetics*, Vol. 35, No. 2, pp. 125-127
- Chan, K. and Kan, Y.W. (1999), "Nrf2 is essential for protection against acute pulmonary injury in mice", *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 96, No. 22, pp. 12731-12736
- Chin, L.S., Olzmann, J.A. and Li, L. (2010), "Parkin-mediated Ubiquitin signalling in aggresome formation and autophagy", *Biochemical Society Transactions*, Vol. 38, No. 1, pp. 144-149
- Choi, P., Golts, N., Snyder, H., Chong, M., Petrucelli, L., Hardy, J., Sparkman, D., Cochran, E., Lee, J.M. and Wolozin, B. (2001), "Co-association of parkin and α -synuclein", *NeuroReport*, Vol. 12, No. 13, pp. 2839-2843
- Chu, J., Hong, N.A., Masuda, C.A., Jenkins, B.V., Nelms, K.A., Goodnow, C.C., Glynn, R.J., Wu, H., Masliah, E., Joazeiro, C.A.P. and Kay, S.A. (2009), "A mouse forward genetics screen identifies LISTERIN as an E3 Ubiquitin ligase involved in neurodegeneration", *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 106, No. 7, pp. 2097-2103
- Ciechanover, A. and Brundin, P. (2003), "The Ubiquitin proteasome system in neurodegenerative diseases: Sometimes the chicken, sometimes the egg", *Neuron*, Vol. 40, No. 2, pp. 427-446
- Cockman, M.E., Masson, N., Mole, D.R., Jaakkola, P., Chang, G.W., Clifford, S.C., Maher, E.R., Pugh, C.W., Ratcliffe, P.J. and Maxwell, P.H. (2000), "Hypoxia inducible factor- α binding and ubiquitylation by the von Hippel-Lindau tumor suppressor protein", *Journal of Biological Chemistry*, Vol. 275, No. 33, pp. 25733-25741

- Collart, M.A. (2003), "Global control of gene expression in yeast by the Ccr4-Not complex", *Gene*, Vol. 313, No. 1-2, pp. 1-16
- Connell, P., Ballinger, C.A., Jiang, J., Wu, Y., Thompson, L.J., Höhfeld, J. and Patterson, C. (2001), "The co-chaperone CHIP regulates protein triage decisions mediated by heat-shock proteins", *Nature Cell Biology*, Vol. 3, No. 1, pp. 93-96
- Cooray, S.N., Guasti, L. and Clark, A.J. (2011), "The E3 Ubiquitin ligase Mahogunin ubiquitinates the melanocortin 2 receptor", *Endocrinology*, Vol. 152, No. 11, pp. 4224-4231
- Cota, C.D., Bagher, P., Pelc, P., Smith, C.O., Bodner, C.R. and Gunn, T.M. (2006), "Mice with mutations in Mahogunin ring finger-1 (*Mgrn1*) exhibit abnormal patterning of the left-right axis", *Developmental Dynamics*, Vol. 235, No. 12, pp. 3438-3447
- Cummings, C.J., Sun, Y., Opal, P., Antalffy, B., Mestril, R., Orr, H.T., Dillmann, W.H. and Zoghbi, H.Y. (2001), "Over-expression of inducible HSP70 chaperone suppresses neuropathology and improves motor function in SCA1 mice", *Human Molecular Genetics*, Vol. 10, No. 14, pp. 1511-1518
- Cyr, D.M., Höhfeld, J. and Patterson, C. (2002), "Protein quality control: U-box-containing E3 Ubiquitin ligases join the fold", *Trends in Biochemical Sciences*, Vol. 27, No. 7, pp. 368-375
- Dai, Q., Zhang, C., Wu, Y., McDonough, H., Whaley, R.A., Godfrey, V., Li, H.H., Madamanchi, N., Xu, W., Neckers, L., Cyr, D. and Patterson, C. (2003), "CHIP activates HSF1 and confers protection against apoptosis and cellular stress", *EMBO Journal*, Vol. 22, No. 20, pp. 5446-5458
- Darom, A., Bening-Abu-Shach, U. and Broday, L. (2010), "RNF-121 is an endoplasmic reticulum-membrane E3 Ubiquitin ligase involved in the regulation of β -integrin", *Molecular Biology of the Cell*, Vol. 21, No. 11, pp. 1788-1798
- Davies, K.J. (1987), "Protein damage and degradation by oxygen radicals. I. general aspects", *Journal of Biological Chemistry*, Vol. 262, No. 20, pp. 9895-9901
- Deak, P.M. and Wolf, D.H. (2001), "Membrane Topology and Function of Der3/Hrd1p as a Ubiquitin-Protein Ligase (E3) Involved in Endoplasmic Reticulum Degradation", *Journal of Biological Chemistry*, Vol. 276, No. 14, pp. 10663-10669
- Delaunay, A., Bromberg, K.D., Hayashi, Y., Mirabella, M., Burch, D., Kirkwood, B., Serra, C., Malicdan, M.C., Mizisin, A.P., Morosetti, R., Broccolini, A., Guo, L.T., Jones, S.N., Lira, S.A., Puri, P.L., Shelton, G.D. and Ronai, Z. (2008), "The ER-bound RING finger protein 5 (RNF5/RMA1) causes degenerative myopathy in transgenic mice and is deregulated in inclusion body myositis", *PLoS ONE*, Vol. 3, No. 2, pp. 1-6
- Denic, V., Quan, E.M. and Weissman, J.S. (2006), "A Luminal Surveillance Complex that Selects Misfolded Glycoproteins for ER-Associated Degradation", *Cell*, Vol. 126, No. 2, pp. 349-359
- Dikshit, P. and Jana, N.R. (2007), "The co-chaperone CHIP is induced in various stresses and confers protection to cells", *Biochemical and Biophysical Research Communications*, Vol. 357, No. 3, pp. 761-765
- Dindot, S.V., Antalffy, B.A., Bhattacharjee, M.B. and Beaudet, A.L. (2008), "The Angelman syndrome Ubiquitin ligase localizes to the synapse and nucleus, and maternal deficiency results in abnormal dendritic spine morphology", *Human Molecular Genetics*, Vol. 17, No. 1, pp. 111-118
- Dobson, C.M. (1999), "Protein misfolding, evolution and disease", *Trends in Biochemical Sciences*, Vol. 24, No. 9, pp. 329-332
- Dobson, C.M. (2003), "Protein folding and misfolding", *Nature*, Vol. 426, No. 6968, pp. 884-890
- Dunah, A.W., Jeong, H., Griffin, A., Kim, Y.M., Standaert, D.G., Hersch, S.M., Mouradian, M.M., Young, A.B., Tanese, N. and Krainc, D. (2002), "Sp1 and TAFII130 transcriptional activity disrupted in early Huntington's disease", *Science*, Vol. 296, No. 5576, pp. 2238-2243
- Egger, L., Madden, D.T., Rhême, C., Rao, R.V. and Bredesen, D.E. (2007), "Endoplasmic reticulum stress-induced cell death mediated by the proteasome", *Cell Death and Differentiation*, Vol. 14, No. 6, pp. 1172-1180
- Ehrlich, E.S., Wang, T., Luo, K., Xiao, Z., Niewiadomska, A.M., Martinez, T., Xu, W., Neckers, L. and Yu, X.F. (2009), "Regulation of Hsp90 client proteins by a Cullin5-RING E3 Ubiquitin ligase", *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 106, No. 48, pp. 20330-20335
- Eisele, F. and Wolf, D.H. (2008), "Degradation of misfolded protein in the cytoplasm is mediated by the Ubiquitin ligase Ubr1", *Federation of European Biochemical Societies Letters*, Vol. 582, No. 30, pp. 4143-4146
- Ellgaard, L., Molinari, M. and Helenius, A. (1999), "Setting the standards: Quality control in the secretory pathway", *Science*, Vol. 286, No. 5446, pp. 1882-1888
- Ellis, J. (1988), "Proteins as molecular chaperones", *Nature*, Vol. 328, No. 6129, pp. 378-379

- Fang, N.N., Ng, A.H.M., Measday, V. and Mayor, T. (2011), "Hul5 HECT Ubiquitin ligase plays a major role in the ubiquitylation and turnover of cytosolic misfolded proteins", *Nature Cell Biology*, Vol. 13, No. 11, pp. 1344-1352
- Fang, S., Ferrone, M., Yang, C., Jensen, J.P., Tiwari, S. and Weissman, A.M. (2001), "The tumor autocrine motility factor receptor, gp78, is a Ubiquitin protein ligase implicated in degradation from the endoplasmic reticulum", *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 98, No. 25, pp. 14422-14427
- Fiedler, K. and Simons, K. (1995), "The role of N-glycans in the secretory pathway", *Cell*, Vol. 81, No. 3, pp. 309-312
- Fu, L., Gao, Y.S. and Sztul, E. (2005), "Transcriptional repression and cell death induced by nuclear aggregates of non-polyglutamine protein", *Neurobiology of Disease*, Vol. 20, No. 3, pp. 656-665
- Ganesh, S., Lal Agarwala, K., Ueda, K., Akagi, T., Shoda, K., Usui, T., Hashikawa, T., Osada, H., Delgado-Escueta, A.V. and Yamakawa, K. (2000), "Laforin, defective in the progressive myoclonus epilepsy of Lafora type, is a dual-specificity phosphatase associated with polyribosomes", *Human Molecular Genetics*, Vol. 9, No. 15, pp. 2251-2261
- Ganesh, S., Puri, R., Singh, S., Mittal, S. and Dubey, D. (2006), "Recent advances in the molecular basis of Lafora's progressive myoclonus epilepsy", *Journal of Human Genetics*, Vol. 51, No. 1, pp. 1-8
- Gao, X. and Hu, H. (2008), "Quality control of the proteins associated with neurodegenerative diseases", *Acta Biochimica et Biophysica Sinica*, Vol. 40, No. 7, pp. 612-618
- Garyali, P., Siwach, P., Singh, P.K., Puri, R., Mittal, S., Sengupta, S., Parihar, R. and Ganesh, S. (2009), "The malin-laforin complex suppresses the cellular toxicity of misfolded proteins by promoting their degradation through the Ubiquitin-proteasome system", *Human Molecular Genetics*, Vol. 18, No. 4, pp. 688-700
- Gentry, M.S., Worby, C.A. and Dixon, J.E. (2005), "Insights into Lafora disease: Malin is an E3 Ubiquitin ligase that ubiquitinates and promotes the degradation of laforin", *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 102, No. 24, pp. 8501-8506
- Glickman, M.H. and Ciechanover, A. (2002), "The Ubiquitin-proteasome proteolytic pathway: Destruction for the sake of construction", *Physiological Reviews*, Vol. 82, No. 2, pp. 373-428
- Goldberg, A.L. (2003), "Protein degradation and protection against misfolded or damaged proteins", *Nature*, Vol. 426, No. 6968, pp. 895-899
- Goswami, A., Dikshit, P., Mishra, A., Nukina, N. and Jana, N.R. (2006), "Expression of expanded polyglutamine proteins suppresses the activation of transcription factor NFkappaB", *Journal of Biological Chemistry*, Vol. 281, No. 48, pp. 37017-37024
- Gourie-Devi, M. (2008), "Organization of neurology services in India: unmet needs and the way forward", *Neurology India*, Vol. 56, No. 1, pp. 4-12
- Greer, P.L., Hanayama, R., Bloodgood, B.L., Mardinly, A.R., Lipton, D.M., Flavell, S.W., Kim, T.K., Griffith, E.C., Waldon, Z., Maehr, R., Ploegh, H.L., Chowdhury, S., Worley, P.F., Steen, J. and Greenberg, M.E. (2010), "The Angelman Syndrome Protein Ube3A Regulates Synapse Development by Ubiquitinating Arc", *Cell*, Vol. 140, No. 5, pp. 704-716
- Groisman, B., Avezov, E. and Lederkremer, G.Z. (2006), "The E3 Ubiquitin ligases HRD1 and SCFFb2 recognize the protein moiety and sugar chains, respectively, of an ER-associated degradation substrate", *Israel Journal of Chemistry*, Vol. 46, No. 2, pp. 189-196
- Grune, T., Jung, T., Merker, K. and Davies, K.J.A. (2004), "Decreased proteolysis caused by protein aggregates, inclusion bodies, plaques, lipofuscin, ceroid, and 'aggresomes' during oxidative stress, aging, and disease", *International Journal of Biochemistry and Cell Biology*, Vol. 36, No. 12, pp. 2519-2530
- Guerra, D.D., Pratelli, R., Kraft, E., Callis, J. and Pilot, G. (2013), "Functional conservation between mammalian MGRN1 and plant LOG2 Ubiquitin ligases", *Federation of European Biochemical Societies Lett*, Vol. 587, No. 21, pp. 3400-3405
- Gunn, T.M., Silvius, D., Bagher, P., Sun, K. and Walker, K.K. (2013), "MGRN1-dependent pigment-type switching requires its ubiquitination activity but not its interaction with TSG101 or NEDD4", *Pigment Cell & Melanoma Research*, Vol. 26, No. 2, pp. 263-268
- Guo, X., Shen, S., Song, S., He, S., Cui, Y., Xing, G., Wang, J., Yin, Y., Fan, L., He, F. and Zhang, L. (2011), "The E3 ligase Smurf1 regulates Wolfram syndrome protein stability at the endoplasmic reticulum", *Journal of Biological Chemistry*, Vol. 286, No. 20, pp. 18037-18047
- Hampton, R.Y. (2002), "ER-associated degradation in protein quality control and cellular regulation", *Current Opinion in Cell Biology*, Vol. 14, No. 4, pp. 476-482

- Hampton, R.Y. (2011), "San1-mediated quality control: Substrate recognition "sans" chaperones", *Molecular Cell*, Vol. 41, No. 1, pp. 2-3
- Harbord, M. (2001), "Levodopa responsive Parkinsonism in adults with Angelman Syndrome", *Journal of Clinical Neuroscience*, Vol. 8, No. 5, pp. 421-422
- Harjes, P. and Wanker, E.E. (2003), "The hunt for huntingtin function: Interaction partners tell many different stories", *Trends in Biochemical Sciences*, Vol. 28, No. 8, pp. 425-433
- Hartl, F.U. and Hayer-Hartl, M. (2002), "Molecular chaperones in the cytosol: From nascent chain to folded protein", *Science*, Vol. 295, No. 5561, pp. 1852-1858
- Hatakeyama, S., Matsumoto, M., Kamura, T., Murayama, M., Chui, D.H., Planel, E., Takahashi, R., Nakayama, K.I. and Takashima, A. (2004), "U-box protein carboxyl terminus of Hsc70-interacting protein (CHIP) mediates poly-ubiquitylation preferentially on four-repeat Tau and is involved in neurodegeneration of tauopathy", *Journal of Neurochemistry*, Vol. 91, No. 2, pp. 299-307
- He, L., Eldridge, A.G., Jackson, P.K., Gunn, T.M. and Barsh, G.S. (2003), "Accessory proteins for melanocortin signaling: Attractin and mahogunin", *Annals of the New York Academy of Sciences*, Vol. 994, pp. 288-298
- He, L., Lu, X.Y., Jolly, A.F., Eldridge, A.G., Watson, S.J., Jackson, P.K., Barsh, G.S. and Gunn, T.M. (2003), "Spongiform degeneration in mahoganoid mutant mice", *Science*, Vol. 299, No. 5607, pp. 710-712
- Heck, J.W., Cheung, S.K. and Hampton, R.Y. (2010), "Cytoplasmic protein quality control degradation mediated by parallel actions of the E3 Ubiquitin ligases Ubr1 and San1", *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 107, No. 3, pp. 1106-1111
- Helenius, A. and Aebi, M. (2001), "Intracellular functions of N-linked glycans", *Science*, Vol. 291, No. 5512, pp. 2364-2369
- Hershko, A. and Ciechanover, A. (1998), "The Ubiquitin system", *Annual Review of Biochemistry*, Vol. 67, pp. 425-479
- Hishikawa, N., Niwa, J.I., Doyu, M., Ito, T., Ishigaki, S., Hashizume, Y. and Sobue, G. (2003), "Dorfin localizes to the ubiquitylated inclusions in Parkinson's disease, dementia with Lewy bodies, multiple system atrophy, and amyotrophic lateral sclerosis", *American Journal of Pathology*, Vol. 163, No. 2, pp. 609-619
- Hoffner, G., Kahlem, P. and Djian, P. (2002), "Perinuclear localization of huntingtin as a consequence of its binding to microtubules through an interaction with beta-tubulin: relevance to Huntington's disease", *Journal of Cell Science*, Vol. 115, No. Pt 5, pp. 941-948
- Hwang, C.S., Shemorry, A., Auerbach, D. and Varshavsky, A. (2010), "The N-end rule pathway is mediated by a complex of the RING-type Ubr1 and HECT-type Ufd4 Ubiquitin ligases", *Nature Cell Biology*, Vol. 12, No. 12, pp. 1177-1185
- Hwang, C.S., Shemorry, A. and Varshavsky, A. (2009), "Two proteolytic pathways regulate DNA repair by cotargeting the Mgt1 alkylguanine transferase", *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 106, No. 7, pp. 2142-2147
- Imai, Y., Soda, M., Hatakeyama, S., Akagi, T., Hashikawa, T., Nakayama, K.I. and Takahashi, R. (2002), "CHIP is associated with Parkin, a gene responsible for familial Parkinson's Disease, and enhances its Ubiquitin ligase activity", *Molecular Cell*, Vol. 10, No. 1, pp. 55-67
- Imai, Y., Soda, M. and Takahashi, R. (2000), "Parkin suppresses unfolded protein stress-induced cell death through its E3 Ubiquitin-protein ligase activity", *Journal of Biological Chemistry*, Vol. 275, No. 46, pp. 35661-35664
- Ishigaki, S., Niwa, J.i., Yamada, S.i., Takahashi, M., Ito, T., Sone, J., Doyu, M., Urano, F. and Sobue, G. (2007), "Dorfin-CHIP chimeric proteins potently ubiquitylate and degrade familial ALS-related mutant SOD1 proteins and reduce their cellular toxicity", *Neurobiology of Disease*, Vol. 25, No. 2, pp. 331-341
- Ismail, N. and Ng, D.T.W. (2006), "Have you HRD? Understanding ERAD Is DOAble!", *Cell*, Vol. 126, No. 2, pp. 237-239
- Iwai, K. (2003), "An Ubiquitin ligase recognizing a protein oxidized by iron: Implications for the turnover of oxidatively damaged proteins", *Journal of Biochemistry*, Vol. 134, No. 2, pp. 175-182
- Iyer, N.V., Kotch, L.E., Agani, F., Leung, S.W., Laughner, E., Wenger, R.H., Gassmann, M., Gearhart, J.D., Lawler, A.M., Yu, A.Y. and Semenza, G.L. (1998), "Cellular and developmental control of O2 homeostasis by hypoxia-inducible factor α ", *Genes and Development*, Vol. 12, No. 2, pp. 149-162
- Jana, N.R., Dikshit, P., Goswami, A., Kotliarova, S., Murata, S., Tanaka, K. and Nukina, N. (2005), "Co-chaperone CHIP associates with expanded polyglutamine protein and promotes their degradation by proteasomes", *Journal of Biological Chemistry*, Vol. 280, No. 12, pp. 11635-11640

- Jiang, H., Ren, Y., Zhao, J. and Feng, J. (2004), "Parkin protects human dopaminergic neuroblastoma cells against dopamine-induced apoptosis", *Human Molecular Genetics*, Vol. 13, No. 16, pp. 1745-1754
- Jiang, J., Ballinger, C.A., Wu, Y., Dai, Q., Cyr, D.M., Höhfeld, J. and Patterson, C. (2001), "CHIP is a U-box-dependent E3 Ubiquitin ligase: Identification of Hsc70 as a target for ubiquitylation", *Journal of Biological Chemistry*, Vol. 276, No. 46, pp. 42938-42944
- Jiao, J., Sun, K., Walker, W.P., Bagher, P., Cota, C.D. and Gunn, T.M. (2009), "Abnormal regulation of TSG101 in mice with spongiform neurodegeneration", *Biochimica et Biophysica Acta*, Vol. 1792, No. 10, pp. 1027-1035
- Johnson, E.S., Ma, P.C.M., Ota, I.M. and Varshavsky, A. (1995), "A proteolytic pathway that recognizes Ubiquitin as a degradation signal", *Journal of Biological Chemistry*, Vol. 270, No. 29, pp. 17442-17456
- Johnston, J.A., Ward, C.L. and Kopito, R.R. (1998), "Aggresomes: A cellular response to misfolded proteins", *Journal of Cell Biology*, Vol. 143, No. 7, pp. 1883-1898
- Junn, E., Lee, S.S., Suhr, U.T. and Mouradian, M.M. (2002), "Parkin accumulation in aggresomes due to proteasome impairment", *Journal of Biological Chemistry*, Vol. 277, No. 49, pp. 47870-47877
- Kahle, P.J. and Haass, C. (2004), "How does parkin ligate Ubiquitin to Parkinson's disease? First in molecular medicine review series", *EMBO Reports*, Vol. 5, No. 7, pp. 681-685
- Kaneko, M., Ishiguro, M., Niinuma, Y., Uesugi, M. and Nomura, Y. (2002), "Human HRD1 protects against ER stress-induced apoptosis through ER-associated degradation", *Federation of European Biochemical Societies Letters*, Vol. 532, No. 1-2, pp. 147-152
- Kawajiri, S., Saiki, S., Sato, S., Sato, F., Hatano, T., Eguchi, H. and Hattori, N. (2010), "PINK1 is recruited to mitochondria with parkin and associates with LC3 in mitophagy", *Federation of European Biochemical Societies Letters*, Vol. 584, No. 6, pp. 1073-1079
- Kikkert, M., Doolman, R., Dai, M., Avner, R., Hassink, G., Van Voorden, S., Thanedar, S., Roitelman, J., Chau, V. and Wiertz, E. (2004), "Human HRD1 Is an E3 Ubiquitin Ligase Involved in Degradation of Proteins from the Endoplasmic Reticulum", *Journal of Biological Chemistry*, Vol. 279, No. 5, pp. 3525-3534
- Kim, B.Y., Olzmann, J.A., Barsh, G.S., Chin, L.S. and Li, L. (2007), "Spongiform neurodegeneration-associated E3 ligase Mahogunin ubiquitylates TSG101 and regulates endosomal trafficking", *Molecular Biology of the Cell*, Vol. 18, No. 4, pp. 1129-1142
- Kishino, T., Lalande, M. and Wagstaff, J. (1997), "UBE3A/E6-AP mutations cause Angelman syndrome", *Nature Genetics*, Vol. 15, No. 1, pp. 70-73
- Kobayashi, A., Kang, M.I., Okawa, H., Ohtsuji, M., Zenke, Y., Chiba, T., Igarashi, K. and Yamamoto, M. (2004), "Oxidative stress sensor Keap1 functions as an adaptor for Cul3-based E3 ligase to regulate proteasomal degradation of Nrf2", *Molecular and Cellular Biology*, Vol. 24, No. 16, pp. 7130-7139
- Koegl, M., Hoppe, T., Schlenker, S., Ulrich, H.D., Mayer, T.U. and Jentsch, S. (1999), "A novel ubiquitination factor, E4, is involved in multiUbiquitin chain assembly", *Cell*, Vol. 96, No. 5, pp. 635-644
- Kon, M. and Cuervo, A.M. (2010), "Autophagy: An alternative degradation mechanism for misfolded proteins", *John Wiley & Sons Incorporation, Hoboken*, pp 113-129
- Kopito, R.R. (2000), "Aggresomes, inclusion bodies and protein aggregation", *Trends in Cell Biology*, Vol. 10, No. 12, pp. 524-530
- Koplin, A., Preissler, S., Llina, Y., Koch, M., Scior, A., Erhardt, M. and Deuerling, E. (2010), "A dual function for chaperones SSB-RAC and the NAC nascent polypeptide-associated complex on ribosomes", *Journal of Cell Biology*, Vol. 189, No. 1, pp. 57-68
- Kopski, K.M. and Huffaker, T.C. (1997), "Suppressors of the ndc10-2 mutation: A role for the Ubiquitin system in *Saccharomyces cerevisiae* kinetochore function", *Genetics*, Vol. 147, No. 2, pp. 409-420
- Korolchuk, V.I., Mansilla, A., Menzies, F.M. and Rubinsztein, D.C. (2009), "Autophagy Inhibition Compromises Degradation of Ubiquitin-Proteasome Pathway Substrates", *Molecular Cell*, Vol. 33, No. 4, pp. 517-527
- Kramer, G., Boehringer, D., Ban, N. and Bukau, B. (2009), "The ribosome as a platform for co-translational processing, folding and targeting of newly synthesized proteins", *Nature Structural and Molecular Biology*, Vol. 16, No. 6, pp. 589-597
- Kreft, S.G., Wang, L. and Hochstrasser, M. (2006), "Membrane topology of the yeast endoplasmic reticulum-localized Ubiquitin ligase Doa10 and comparison with its human ortholog TEB4 (MARCH-VI)", *Journal of Biological Chemistry*, Vol. 281, No. 8, pp. 4646-4653

- Krüger, R., Kuhn, W., Müller, T., Voitalla, D., Graeber, M., Kösel, S., Przuntek, H., Epplen, J.T., Schols, L. and Riess, O. (1998), "Ala30Pro mutation in the gene encoding α -synuclein in Parkinson's disease", *Nature Genetics*, Vol. 18, No. 2, pp. 106-108
- Kubota, H. (2009), "Quality control against misfolded proteins in the cytosol: A network for cell survival", *Journal of Biochemistry*, Vol. 146, No. 5, pp. 609-616
- Kumar, P., Ambasta, R.K., Veereshwarayya, V., Rosen, K.M., Kosik, K.S., Band, H., Mestril, R., Patterson, C. and Querfurth, H.W. (2007), "CHIP and HSPs interact with β -APP in a proteasome-dependent manner and influence A β metabolism", *Human Molecular Genetics*, Vol. 16, No. 7, pp. 848-864
- Kwon, Y.T., Reiss, Y., Fried, V.A., Hershko, A., Yoon, J.K., Gonda, D.K., Sangan, P., Copeland, N.G., Jenkins, N.A. and Varshavsky, A. (1998), "The mouse and human genes encoding the recognition component of the N-end rule pathway", *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 95, No. 14, pp. 7898-7903
- Lang, A.E. and Lozano, A.M. (1998), "Parkinson's disease", *New England Journal of Medicine*, Vol. 339, No. 15, pp. 1044-1053
- Lee, D.H., Sherman, M.Y. and Goldberg, A.L. (1996), "Involvement of the molecular chaperone Ydj1 in the Ubiquitin-dependent degradation of short-lived and abnormal proteins in *Saccharomyces cerevisiae*", *Molecular and Cellular Biology*, Vol. 16, No. 9, pp. 4773-4781
- Leiser, S.F. and Miller, R.A. (2010), "Nrf2 signaling, a mechanism for cellular stress resistance in long-lived mice", *Molecular and Cellular Biology*, Vol. 30, No. 3, pp. 871-884
- Lenartowski, R., Gumowski, K. and Goc, A. (2008), "The multifunctionality of CHIP protein in the protein quality-control system", *Postępy higieny i medycyny doświadczalnej*, Vol. 62, No., pp. 297-308
- Lerner, M., Corcoran, M., Cepeda, D., Nielsen, M.L., Zubarev, R., Pontén, F., Uhlén, M., Hober, S., Grandér, D. and Sangfelt, O. (2007), "The RBCC gene RFP2 (Leu5) encodes a novel transmembrane E3 Ubiquitin ligase involved in ERAD", *Molecular Biology of the Cell*, Vol. 18, No. 5, pp. 1670-1682
- Levine, B. and Kroemer, G. (2008), "Autophagy in the Pathogenesis of Disease", *Cell*, Vol. 132, No. 1, pp. 27-42
- Levy, F., Muehlethaler, K., Salvi, S., Peitrequin, A.L., Lindholm, C.K., Cerottini, J.C. and Rimoldi, D. (2005), "Ubiquitylation of a melanosomal protein by HECT-E3 ligases serves as sorting signal for lysosomal degradation", *Molecular Biology of the Cell*, Vol. 16, No. 4, pp. 1777-1787
- Li, J.Y., Englund, E., Widner, H., Rehncrona, S., Björklund, A., Lindvall, O. and Brundin, P. (2010), "Characterization of Lewy body pathology in 12- and 16-year-old intrastriatal mesencephalic grafts surviving in a patient with Parkinson's disease", *Movement Disorders*, Vol. 25, No. 8, pp. 1091-1096
- Li, S.H., Cheng, A.L., Zhou, H., Lam, S., Rao, M., Li, H. and Li, X.J. (2002), "Interaction of Huntington disease protein with transcriptional activator Sp1", *Mol Cell Biol*, Vol. 22, No. 5, pp. 1277-1287
- Lilley, B.N. and Ploegh, H.L. (2004), "A membrane protein required for dislocation of misfolded proteins from the ER", *Nature*, Vol. 429, No. 6994, pp. 834-840
- Löffek, S., Wöll, S., Höhfeld, J., Leube, R.E., Has, C., Bruckner-Tuderman, L. and Magin, T.M. (2010), "The Ubiquitin ligase CHIP/STUB1 targets mutant keratins for degradation", *Human Mutation*, Vol. 31, No. 4, pp. 466-476
- Lowe, J., Blanchard, A., Morrell, K., Lennox, G., Reynolds, L., Billett, M., Landon, M. and Mayer, R.J. (1988), "Ubiquitin is a common factor in intermediate filament inclusion bodies of diverse type in man, including those of Parkinson's disease, Pick's disease, and Alzheimer's disease, as well as Rosenthal fibres in cerebellar astrocytomas, cytoplasmic bodies in muscle, and Mallory bodies in alcoholic liver disease", *Journal of Pathology*, Vol. 155, No. 1, pp. 9-15
- Maeda, T., Marutani, T., Zou, K., Araki, W., Tanabe, C., Yagishita, N., Yamano, Y., Amano, T., Michikawa, M., Nakajima, T. and Komano, H. (2009), "An E3 Ubiquitin ligase, Synoviolin, is involved in the degradation of immature nicastrin, and regulates the production of amyloid β -protein", *Federation of European Biochemical Societies Journal*, Vol. 276, No. 20, pp. 5832-5840
- Mangiarini, L., Sathasivam, K., Seller, M., Cozens, B., Harper, A., Hetherington, C., Lawton, M., Trotter, Y., Lehrach, H., Davies, S.W. and Bates, G.P. (1996), "Exon 1 of the HD gene with an expanded CAG repeat is sufficient to cause a progressive neurological phenotype in transgenic mice", *Cell*, Vol. 87, No. 3, pp. 493-506
- Margolis, R.L. and Ross, C.A. (2001), "Expansion explosion: new clues to the pathogenesis of repeat expansion neurodegenerative diseases", *Trends Mol Med*, Vol. 7, No. 11, pp. 479-482
- Martin, J.B. (1999), "Molecular basis of the neurodegenerative disorders", *New England Journal of Medicine*, Vol. 340, No. 25, pp. 1970-1980

- Martinez, V.D., Vucic, E.A., Thu, K.L., Pikor, L.A., Lam, S. and Lam, W.L. (2014), "Disruption of KEAP1/CUL3/RBX1 E3-Ubiquitin ligase complex components by multiple genetic mechanisms is associated with poor prognosis in head and neck cancer", *Journal of Sciences and Specialities of The Head and Neck*, Head & Neck, DOI:10.1002/hed.23663
- Maruyama, Y., Yamada, M. and Takahashi, K. (2008), "Ubiquitin ligase Kf-1 is involved in the endoplasmic reticulum-associated degradation pathway", *Biochemical and Biophysical Research Communications*, Vol. 374, No. 4, pp. 737-741
- Massey, A.C., Zhang, C. and Cuervo, A.M. (2006), "Chaperone-Mediated Autophagy in Aging and Disease", *Current Topics in Developmental Biology*, Vol. 73, pp. 205-235
- Matsuo, Y., Kishimoto, H., Tanae, K., Kitamura, K., Katayama, S. and Kawamukai, M. (2011), "Nuclear protein quality is regulated by the Ubiquitin-proteasome system through the activity of Ubc4 and San1 in fission yeast", *Journal of Biological Chemistry*, Vol. 286, No. 15, pp. 13775-13790
- Maynard, C.J., Bottcher, C., Ortega, Z., Smith, R., Florea, B.I., Diaz-Hernandez, M., Brundin, P., Overkleeft, H.S., Li, J.Y., Lucas, J.J. and Dantuma, N.P. (2009), "Accumulation of Ubiquitin conjugates in a polyglutamine disease model occurs without global ubiquitin/proteasome system impairment", *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 106, No. 33, pp. 13986-13991
- McClellan, A.J. and Frydman, J. (2001), "Molecular chaperones and the art of recognizing a lost cause", *Nature Cell Biology*, Vol. 3, No. 2, pp. E51-E53
- McClellan, A.J., Scott, M.D. and Frydman, J. (2005), "Folding and quality control of the VHL tumor suppressor proceed through distinct chaperone pathways", *Cell*, Vol. 121, No. 5, pp. 739-748
- Meacham, G.C., Patterson, C., Zhang, W., Younger, J.M. and Cyr, D.M. (2001), "The Hsc70 co-chaperone CHIP targets immature CFTR for proteasomal degradation", *Nature Cell Biology*, Vol. 3, No. 1, pp. 100-105
- Menzies, F.M., Moreau, K. and Rubinsztein, D.C. (2011), "Protein misfolding disorders and macroautophagy", *Current Opinion in Cell Biology*, Vol. 23, No. 2, pp. 190-197
- Menzies, F.M., Yenissetti, S.C. and Min, K.T. (2005), "Roles of Drosophila DJ-1 in survival of dopaminergic neurons and oxidative stress", *Current Biology*, Vol. 15, No. 17, pp. 1578-1582
- Meulener, M., Whitworth, A.J., Armstrong-Gold, C.E., Rizzu, P., Heutink, P., Wes, P.D., Pallanck, L.J. and Bonini, N.M. (2005), "Drosophila DJ-1 mutants are selectively sensitive to environmental toxins associated with Parkinson's disease", *Current Biology*, Vol. 15, No. 17, pp. 1572-1577
- Miller, K.A., Gunn, T.M., Carrasquillo, M.M., Lamoreux, M.L., Galbraith, D.B. and Barsh, G.S. (1997), "Genetic studies of the mouse mutations mahogany and mahoganoid", *Genetics*, Vol. 146, No. 4, pp. 1407-1415
- Miller, V.M., Nelson, R.F., Gouvion, C.M., Williams, A., Rodriguez-Lebron, E., Harper, S.Q., Davidson, B.L., Rebagliati, M.R. and Paulson, H.L. (2005), "CHIP suppresses polyglutamine aggregation and toxicity in vitro and in vivo", *Journal of Neuroscience*, Vol. 25, No. 40, pp. 9152-9161
- Minassian, B.A., Lee Jeffrey, R., Herbrick, J.A., Huizenga, J., Soder, S., Mungall, A.J., Dunham, I., Gardner, R., Fong, C.Y.G., Carpenter, S., Jardim, L., Satishchandra, P., Andermann, E., Carter Snead Iii, O., Lopes-Cendes, I., Tsui, L.C., Delgado-Escueta, A.V., Rouleau, G.A. and Scherer, S.W. (1998), "Mutations in a gene encoding a novel protein tyrosine phosphatase cause progressive myoclonus epilepsy", *Nature Genetics*, Vol. 20, No. 2, pp. 171-174
- Mishra, A. (2013), "E6-AP association promotes SOD1 aggresomes degradation and suppresses toxicity", *Neurobiology of Aging*, Vol. 34, No. 4, pp.1310.e11-23
- Mishra, A., Dikshit, P., Purkayastha, S., Sharma, J., Nukina, N. and Jana, N.R. (2008), "E6-AP promotes misfolded polyglutamine proteins for proteasomal degradation and suppresses polyglutamine protein aggregation and toxicity", *Journal of Biological Chemistry*, Vol. 283, No. 12, pp. 7648-7656
- Mishra, A., Godavarthi, S.K. and Jana, N.R. (2009), "UBE3A/E6-AP regulates cell proliferation by promoting proteasomal degradation of p27", *Neurobiology of Disease*, Vol. 36, No. 1, pp. 26-34
- Mishra, A., Godavarthi, S.K., Maheshwari, M., Goswami, A. and Jana, N.R. (2009), "The Ubiquitin ligase E6-AP is induced and recruited to aggresomes in response to proteasome inhibition and may be involved in the ubiquitination of Hsp70-bound misfolded proteins", *Journal of Biological Chemistry*, Vol. 284, No. 16, pp. 10537-10545
- Mishra, A. and Jana, N.R. (2008), "Regulation of turnover of tumor suppressor p53 and cell growth by E6-AP, a Ubiquitin protein ligase mutated in Angelman mental retardation syndrome", *Cellular and Molecular Life Sciences*, Vol. 65, No. 4, pp. 656-666

- Mitsui, K., Nakayama, H., Akagi, T., Nekooki, M., Ohtawa, K., Takio, K., Hashikawa, T. and Nukina, N. (2002), "Purification of polyglutamine aggregates and identification of elongation factor-1 α and heat shock protein 84 as aggregate-interacting proteins", *Journal of Neuroscience*, Vol. 22, No. 21, pp. 9267-9277
- Mitsumoto, A., Nakagawa, Y., Takeuchi, A., Okawa, K., Iwamatsu, A. and Takanezawa, Y. (2001), "Oxidized forms of peroxiredoxins and DJ-1 on two-dimensional gels increased in response to sublethal levels of paraquat", *Free Radical Research*, Vol. 35, No. 3, pp. 301-310
- Mittal, S., Dubey, D., Yamakawa, K. and Ganesh, S. (2007), "Lafora disease proteins malin and laforin are recruited to aggresomes in response to proteasomal impairment", *Human Molecular Genetics*, Vol. 16, No. 7, pp. 753-762
- Miyazaki, K., Fujita, T., Ozaki, T., Kato, C., Kurose, Y., Sakamoto, M., Kato, S., Goto, T., Itoyama, Y., Aoki, M. and Nakagawara, A. (2004), "NEDL1, a Novel Ubiquitin-protein Isopeptide Ligase for Dishevelled-1, Targets Mutant Superoxide Dismutase-1", *Journal of Biological Chemistry*, Vol. 279, No. 12, pp. 11327-11335
- Mizuno, Y., Hattori, N., Mori, H., Suzuki, T. and Tanaka, K. (2001), "Parkin and Parkinson's disease", *Current Opinion in Neurology*, Vol. 14, No. 4, pp. 477-482
- Mizushima, N., Levine, B., Cuervo, A.M. and Klionsky, D.J. (2008), "Autophagy fights disease through cellular self-digestion", *Nature*, Vol. 451, No. 7182, pp. 1069-1075
- Morito, D., Hirao, K., Oda, Y., Hosokawa, N., Tokunaga, F., Cyr, D.M., Tanaka, K., Iwai, K. and Nagata, K. (2008), "Gp78 cooperates with RMA1 in endoplasmic reticulum-associated degradation of CFTR^{R508}", *Molecular Biology of the Cell*, Vol. 19, No. 4, pp. 1328-1336
- Mulherkar, S.A., Sharma, J. and Jana, N.R. (2009), "The Ubiquitin ligase E6-AP promotes degradation of α -synuclein", *Journal of Neurochemistry*, Vol. 110, No. 6, pp. 1955-1964
- Muller-Reichert, T., Sassoon, I., O'Toole, E., Romao, M., Ashford, A.J., Hyman, A.A. and Antony, C. (2003), "Analysis of the distribution of the kinetochore protein Ndc10p in *Saccharomyces cerevisiae* using 3-D modeling of mitotic spindles", *Chromosoma*, Vol. 111, No. 7, pp. 417-428
- Muqit, M.M.K., Davidson, S.M., Payne Smith, M.D., MacCormac, L.P., Kahns, S., Jensen, P.H., Wood, N.W. and Latchman, D.S. (2004), "Parkin is recruited into aggresomes in a stress-specific manner: Over-expression of parkin reduces aggresome formation but can be dissociated from parkin's effect on neuronal survival", *Human Molecular Genetics*, Vol. 13, No. 1, pp. 117-135
- Murata, S., Minami, Y., Minami, M., Chiba, T. and Tanaka, K. (2001), "CHIP is a chaperone-dependent E3 ligase that ubiquitylates unfolded protein", *EMBO Reports*, Vol. 2, No. 12, pp. 1133-1138
- Nabi, I.R. and Raz, A. (1987), "Cell shape modulation alters glycosylation of a metastatic melanoma cell-surface antigen", *International Journal of Cancer*, Vol. 40, No. 3, pp. 396-402
- Nabi, I.R., Watanabe, H., Silletti, S. and Raz, A. (1991), "Tumor cell autocrine motility factor receptor", *Experientia Supplementum*, Vol. 59, No., pp. 163-177
- Nagase, T., Ishikawa, K., Miyajima, N., Tanaka, A., Kotani, H., Nomura, N. and Ohara, O. (1998), "Prediction of the coding sequences of unidentified human genes. IX. The complete sequences of 100 new cDNA clones from brain which can code for large proteins in vitro", *DNA Research*, Vol. 5, No. 1, pp. 31-39
- Nakamura, T. and Lipton, S.A. (2009), "Cell death: Protein misfolding and neurodegenerative diseases", *Apoptosis*, Vol. 14, No. 4, pp. 455-468
- Nasir, J., Floresco, S.B., O'Kusky, J.R., Diewert, V.M., Richman, J.M., Zeisler, J., Borowski, A., Marth, J.D., Phillips, A.G. and Hayden, M.R. (1995), "Targeted disruption of the Huntington's disease gene results in embryonic lethality and behavioral and morphological changes in heterozygotes", *Cell*, Vol. 81, No. 5, pp. 811-823
- Nelson, R.F., Glenn, K.A., Zhang, Y., Wen, H., Knutson, T., Gouvion, C.M., Robinson, B.K., Zhou, Z., Yang, B., Smith, R.J.H. and Paulson, H.L. (2007), "Selective cochlear degeneration in mice lacking the F-box protein, Fbx2, a glycoprotein-specific Ubiquitin ligase subunit", *Journal of Neuroscience*, Vol. 27, No. 19, pp. 5163-5171
- Nguyen, T., Nioi, P. and Pickett, C.B. (2009), "The Nrf2-antioxidant response element signaling pathway and its activation by oxidative stress", *Journal of Biological Chemistry*, Vol. 284, No. 20, pp. 13291-13295
- Nillegoda, N.B., Theodoraki, M.A., Mandal, A.K., Mayo, K.J., Ren, H.Y., Sultana, R., Wu, K., Johnson, J., Cyr, D.M. and Caplan, A.J. (2010), "Ubr1 and Ubr2 function in a quality control pathway for degradation of unfolded cytosolic proteins", *Molecular Biology of the Cell*, Vol. 21, No. 13, pp. 2102-2116

- Nishitani, H., Sugimoto, N., Roukos, V., Nakanishi, Y., Saijo, M., Obuse, C., Tsurimoto, T., Nakayama, K.I., Nakayama, K., Fujita, M., Lygerou, Z. and Nishimoto, T. (2006), "Two E3 Ubiquitin ligases, SCF-Skp2 and DDB1-Cul4, target human Cdt1 for proteolysis", *EMBO Journal*, Vol. 25, No. 5, pp. 1126-1136
- Niwa, J.I., Ishigaki, S., Doyu, M., Suzuki, T., Tanaka, K. and Sobue, G. (2001), "A novel centrosomal RING-finger protein, Dorfin, mediates Ubiquitin ligase activity", *Biochemical and Biophysical Research Communications*, Vol. 281, No. 3, pp. 706-713
- Niwa, J.I., Ishigaki, S., Hishikawa, N., Yamamoto, M., Doyu, M., Murata, S., Tanaka, K., Taniguchi, N. and Sobue, G. (2002), "Dorfin ubiquitylates mutant SOD1 and prevents mutant SOD1-mediated neurotoxicity", *Journal of Biological Chemistry*, Vol. 277, No. 39, pp. 36793-36798
- Olzmann, J.A., Li, L. and Chin, L.S. (2008), "Aggresome formation and neurodegenerative diseases: Therapeutic implications", *Current Medicinal Chemistry*, Vol. 15, No. 1, pp. 47-60
- Olzsha, H., Schermann, S.M., Woerner, A.C., Pinkert, S., Hecht, M.H., Tartaglia, G.G., Vendruscolo, M., Hayer-Hartl, M., Hartl, F.U. and Vabulas, R.M. (2011), "Amyloid-like aggregates sequester numerous metastable proteins with essential cellular functions", *Cell*, Vol. 144, No. 1, pp. 67-78
- Omura, T., Kaneko, M., Okuma, Y., Orba, Y., Nagashima, K., Takahashi, R., Fujitani, N., Matsumura, S., Hata, A., Kubota, K., Murahashi, K., Uehara, T. and Nomura, Y. (2006), "A Ubiquitin ligase HRD1 promotes the degradation of Pael receptor, a substrate of Parkin", *Journal of Neurochemistry*, Vol. 99, No. 6, pp. 1456-1469
- Ortega, Z., Diaz-Hernandez, M. and Lucas, J.J. (2007), "Is the Ubiquitin-proteasome system impaired in Huntington's disease?", *Cellular and Molecular Life Sciences*, Vol. 64, No. 17, pp. 2245-2257
- Ortega, Z., Diaz-Hernandez, M., Maynard, C.J., Hernandez, F., Dantuma, N.P. and Lucas, J.J. (2010), "Acute polyglutamine expression in inducible mouse model unravels ubiquitin/proteasome system impairment and permanent recovery attributable to aggregate formation", *Journal of Neuroscience*, Vol. 30, No. 10, pp. 3675-3688
- Overton, J.D. and Leibel, R.L. (2011), "Mahoganoid and mahogany mutations rectify the obesity of the yellow mouse by effects on endosomal traffic of MC4R protein", *Journal of Biological Chemistry*, Vol. 286, No. 21, pp. 18914-18929
- Panasenko, O.O. and Collart, M.A. (2012), "Presence of Not5 and ubiquitinated Rps7A in polysome fractions depends upon the Not4 E3 ligase", *Molecular Microbiology*, Vol. 83, No. 3, pp. 640-653
- Pardo, R., Molina-Calavita, M., Poizat, G., Keryer, G., Humbert, S. and Saudou, F. (2010), "pARIS-htt: an optimised expression platform to study huntingtin reveals functional domains required for vesicular trafficking", *Molecular Brain*, Vol. 3, No., pp. 17
- Park, J., Lee, S.B., Lee, S., Kim, Y., Song, S., Kim, S., Bae, E., Kim, J., Shong, M., Kim, J.M. and Chung, J. (2006), "Mitochondrial dysfunction in Drosophila PINK1 mutants is complemented by parkin", *Nature*, Vol. 441, No. 7097, pp. 1157-1161
- Paschen, W. and Frandsen, A. (2001), "Endoplasmic reticulum dysfunction - A common denominator for cell injury in acute and degenerative diseases of the brain?", *Journal of Neurochemistry*, Vol. 79, No. 4, pp. 719-725
- Paul, S. (2008), "Dysfunction of the Ubiquitin-proteasome system in multiple disease conditions: Therapeutic approaches", *BioEssays*, Vol. 30, No. 11-12, pp. 1172-1184
- Perez-Oliva, A.B., Olivares, C., Jimenez-Cervantes, C. and Garcia-Borron, J. (2009), "Mahogunin ring finger-1 (MGRN1) E3 Ubiquitin ligase inhibits signaling from melanocortin receptor by competition with G-alphas", *Journal of Biological Chemistry*, Vol. 284, No. 46, pp. 31714-31725
- Phan, L.K., Lin, F., LeDuc, C.A., Chung, W.K. and Leibel, R.L. (2002), "The mouse mahoganoid coat color mutation disrupts a novel C3HC4 RING domain protein", *J Clin Invest*, Vol. 110, No. 10, pp. 1449-1459
- Pickart, C.M. (2001). "Mechanisms underlying ubiquitination", *Annual Review of Biochemistry*, Vol. 70, pp. 503-533
- Plempner, R.K., Böhmeler, S., Bordallo, J., Sommer, T. and Wolf, D.H. (1997), "Mutant analysis links the translocon and BIP to retrograde protein transport for ER degradation", *Nature*, Vol. 388, No. 6645, pp. 891-895
- Polymeropoulos, M.H., Lavedan, C., Leroy, E., Ide, S.E., Dehejia, A., Dutra, A., Pike, B., Root, H., Rubenstein, J., Boyer, R., Stenroos, E.S., Chandrasekharappa, S., Athanassiadou, A., Papapetropoulos, T., Johnson, W.G., Lazzarini, A.M., Duvoisin, R.C., Di Iorio, G., Golbe, L.I. and Nussbaum, R.L. (1997), "Mutation in the α -synuclein gene identified in families with Parkinson's disease", *Science*, Vol. 276, No. 5321, pp. 2045-2047
- Powers, E.T., Morimoto, R.I., Dillin, A., Kelly, J.W. and Balch, W.E. (2009), "Biological and chemical approaches to diseases of proteostasis deficiency", *Annual Review of Biochemistry*, Vol. 78, pp. 959-991

- Ramaswamy, S., McBride, J.L. and Kordower, J.H. (2007), "Animal models of Huntington's disease", *Institute for Laboratory Animal Research Journal*, Vol. 48, No. 4, pp. 356-373
- Rao, R.V., Ellerby, H.M. and Bredesen, D.E. (2004), "Coupling endoplasmic reticulum stress to the cell death program", *Cell Death and Differentiation*, Vol. 11, No. 4, pp. 372-380
- Rao, S.N., Maity, R., Sharma, J., Dey, P., Shankar, S.K., Satishchandra, P. and Jana, N.R. (2010), "Sequestration of chaperones and proteasome into Lafora bodies and proteasomal dysfunction induced by Lafora disease-associated mutations of malin", *Human Molecular Genetics*, Vol. 19, No. 23, pp. 4726-4734
- Rauch, T., Hundley, H.A., Pfund, C., Wegrzyn, R.D., Walter, W., Kramer, G., Kim, S.Y., Craig, E.A. and Deuerling, E. (2005), "Dissecting functional similarities of ribosome-associated chaperones from *Saccharomyces cerevisiae* and *Escherichia coli*", *Molecular Microbiology*, Vol. 57, No. 2, pp. 357-365
- Ravid, T., Kreft, S.G. and Hochstrasser, M. (2006), "Membrane and soluble substrates of the Doa10 Ubiquitin ligase are degraded by distinct pathways", *EMBO Journal*, Vol. 25, No. 3, pp. 533-543
- Ravikumar, B. and Rubinsztein, D.C. (2006), "Role of autophagy in the clearance of mutant huntingtin: a step towards therapy?", *Molecular Aspects of Medicine*, Vol. 27, No. 5-6, pp. 520-527
- Reggiori, F. and Pelham, H.R.B. (2002), "A transmembrane Ubiquitin ligase required to sort membrane proteins into multivesicular bodies", *Nature Cell Biology*, Vol. 4, No. 2, pp. 117-123
- Reinstein, E. and Ciechanover, A. (2006), "Narrative review: Protein degradation and human diseases: The Ubiquitin connection", *Annals of Internal Medicine*, Vol. 145, No. 9, pp. 676-684
- Ren, Y., Zhao, J. and Feng, J. (2003), "Parkin binds to α/β tubulin and increases their ubiquitination and degradation", *Journal of Neuroscience*, Vol. 23, No. 8, pp. 3316-3324
- Rong, J., Chen, L., Toth, J.I., Tcherpakov, M., Petroski, M.D. and Reed, J.C. (2011), "Bifunctional Apoptosis Regulator (BAR), an Endoplasmic Reticulum (ER)-associated E3 Ubiquitin ligase, modulates BI-1 protein stability and function in ER stress", *Journal of Biological Chemistry*, Vol. 286, No. 2, pp. 1453-1463
- Rosenbaum, J.C., Fredrickson, E.K., Oeser, M.L., Garrett-Engele, C.M., Locke, M.N., Richardson, L.A., Nelson, Z.W., Hetrick, E.D., Milac, T.I., Gottschling, D.E. and Gardner, R.G. (2011), "Disorder targets misorder in nuclear quality control degradation: A disordered Ubiquitin ligase directly recognizes its misfolded substrates", *Molecular Cell*, Vol. 41, No. 1, pp. 93-106
- Ross, C.A. and Poirier, M.A. (2004), "Protein aggregation and neurodegenerative disease", *Nature Medicine*, Vol. 10, SUPPL., pp. S10-S17
- Roth, W., Kermer, P., Krajewska, M., Welsh, K., Davis, S., Krajewski, S. and Reed, J.C. (2003), "Bifunctional apoptosis inhibitor (BAR) protects neurons from diverse cell death pathways", *Cell Death and Differentiation*, Vol. 10, No. 10, pp. 1178-1187
- Schaffar, G., Breuer, P., Boteva, R., Behrends, C., Tzvetkov, N., Strippel, N., Sakahira, H., Siegers, K., Hayer-Hartl, M. and Hartl, F.U. (2004), "Cellular toxicity of polyglutamine expansion proteins: mechanism of transcription factor deactivation", *Molecular Cell*, Vol. 15, No. 1, pp. 95-105
- Schipper-Krom, S., Juenemann, K. and Reits, E.A. (2012), "The Ubiquitin-Proteasome System in Huntington's Disease: Are Proteasomes Impaired, Initiators of Disease, or Coming to the Rescue?", *Biochemistry Research International*, Vol. 2012, pp. 837015(12 pages)
- Schubert, U., Antón, L.C., Gibbs, J., Norbury, C.C., Yewdell, J.W. and Bennink, J.R. (2000), "Rapid degradation of a large fraction of newly synthesized proteins by proteasomes", *Nature*, Vol. 404, No. 6779, pp. 770-774
- Selkoe, D.J. (2003), "Folding proteins in fatal ways", *Nature*, Vol. 426, No. 6968, pp. 900-904
- Seo, K., Mohanty, T.R., Choi, T. and Hwang, I. (2007), "Biology of epidermal and hair pigmentation in cattle: a mini-review", *Veterinary Dermatology*, Vol. 18, No. 6, pp. 392-400
- Serratos, J.M., Gómez-Garre, P., Gallardo, M.E., Anta, B., Beltrán-Valero de Bernabé, D., Lindhout, D., Augustijn, P.B., Tassinari, C.A., Michelucci, R., Malafosse, A., Topcu, M., Grid, D., Dravet, C., Berkovic, S.F. and Rodríguez de Córdoba, S. (1999), "A novel protein tyrosine phosphatase gene is mutated in progressive myoclonus epilepsy of the Lafora type (EPM2)", *Human Molecular Genetics*, Vol. 8, No. 2, pp. 345-352
- Sha, Y., Pandit, L., Zeng, S. and Eissa, N.T. (2009), "A critical role for CHIP in the aggresome pathway", *Molecular and Cellular Biology*, Vol. 29, No. 1, pp. 116-128
- Shah, I.M. and Di Napoli, M. (2007), "The Ubiquitin-proteasome system and proteasome inhibitors in central nervous system diseases", *Cardiovascular and Hematological Disorders - Drug Targets*, Vol. 7, No. 4, pp. 250-273

- Shang, F., Gong, X. and Taylor, A. (1997), "Activity of Ubiquitin-dependent pathway in response to oxidative stress: Ubiquitin-activating enzyme is transiently up-regulated", *Journal of Biological Chemistry*, Vol. 272, No. 37, pp. 23086-23093
- Shen, Y., Ballar, P. and Fang, S. (2006), "Ubiquitin ligase gp78 increases solubility and facilitates degradation of the Z variant of α 1-antitrypsin", *Biochemical and Biophysical Research Communications*, Vol. 349, No. 4, pp. 1285-1293
- Shimohata, T., Onodera, O. and Tsuji, S. (2000), "Interaction of expanded polyglutamine stretches with nuclear transcription factors leads to aberrant transcriptional regulation in polyglutamine diseases", *Neuropathology*, Vol. 20, No. 4, pp. 326-333
- Shimura, H., Hattori, N., Kubo, S.I., Mizuno, Y., Asakawa, S., Minoshima, S., Shimizu, N., Iwai, K., Chiba, T., Tanaka, K. and Suzuki, T. (2000), "Familial Parkinson disease gene product, parkin, is a Ubiquitin-protein ligase", *Nature Genetics*, Vol. 25, No. 3, pp. 302-305
- Shimura, H., Schwartz, D., Gygi, S.P. and Kosik, K.S. (2004), "CHIP-Hsc70 Complex Ubiquitinates Phosphorylated Tau and Enhances Cell Survival", *Journal of Biological Chemistry*, Vol. 279, No. 6, pp. 4869-4876
- Singh, S., Satishchandra, P., Shankar, S.K. and Ganesh, S. (2008), "Lafora disease in the Indian population: EPM2A and NHLRC1 gene mutations and their impact on subcellular localization of laforin and malin", *Human Mutation*, Vol. 29, No. 6, pp. E1-12
- Sisoula, C. and Gonos, E.S. (2011), "CHIP E3 ligase regulates mammalian senescence by modulating the levels of oxidized proteins", *Mechanisms of Ageing and Development*, Vol. 132, No. 5, pp. 269-272
- Song, B.L., Sever, N. and DeBose-Boyd, R.A. (2005), "Gp78, a membrane-anchored Ubiquitin ligase, associates with Insig-1 and couples sterol-regulated ubiquitination to degradation of HMG CoA reductase", *Molecular Cell*, Vol. 19, No. 6, pp. 829-840
- Spillantini, M.G., Crowther, R.A., Jakes, R., Hasegawa, M. and Goedert, M. (1998), "alpha Synuclein in filamentous inclusions of Lewy bodies from Parkinson's disease and dementia with Lewy bodies", *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 95, No. 11, pp. 6469-6473
- Squier, T.C. (2001), "Oxidative stress and protein aggregation during biological aging", *Experimental Gerontology*, Vol. 36, No. 9, pp. 1539-1550
- Srivastava, D. and Chakrabarti, O. (2014), "Mahogunin-mediated alpha-tubulin ubiquitination via noncanonical K6 linkage regulates microtubule stability and mitotic spindle orientation", *Cell Death and Disease*, Vol. 5, No., pp. e1064
- Stadtman, E.R. (1992), "Protein oxidation and aging", *Science*, Vol. 257, No. 5074, pp. 1220-1224
- Stadtman, E.R. and Levine, R.L. (2000), "Protein oxidation", *Annals of the New York Academy of Sciences*, Vol. 899, pp. 191-208
- Starke-Reed, P.E. and Oliver, C.N. (1989), "Protein oxidation and proteolysis during aging and oxidative stress", *Archives of Biochemistry and Biophysics*, Vol. 275, No. 2, pp. 559-567
- Stefanis, L., Kholodilov, N., Rideout, H.J., Burke, R.E. and Greene, L.A. (2001), "Synuclein-1 is selectively up-regulated in response to nerve growth factor treatment in PC12 cells", *Journal of Neurochemistry*, Vol. 76, No. 4, pp. 1165-1176
- Stegmuller, J. and Bonni, A. (2010), "Destroy to create: E3 Ubiquitin ligases in neurogenesis", *F1000 Biology Reports*, Vol. 2, No. 1,
- Sullivan, E.K., Weirich, C.S., Guyon, J.R., Sif, S. and Kingston, R.E. (2001), "Transcriptional activation domains of human heat shock factor 1 recruit human SWI/SNF", *Molecular and Cellular Biology*, Vol. 21, No. 17, pp. 5826-5837
- Sun, K., Johnson, B.S. and Gunn, T.M. (2007), "Mitochondrial dysfunction precedes neurodegeneration in mahogunin (Mgn1) mutant mice", *Neurobiology of Aging*, Vol. 28, No. 12, pp. 1840-1852
- Taira, T., Saito, Y., Niki, T., Iguchi-Arigo, S.M.M., Takahashi, K. and Ariga, H. (2004), "DJ-1 has a role in antioxidative stress to prevent cell death", *EMBO Reports*, Vol. 5, No. 2, pp. 213-218
- Tanaka, Y., Engelender, S., Igarashi, S., Rao, R.K., Wanner, T., Tanzi, R.E., Sawa, A., Dawson, V.L., Dawson, T.M. and Ross, C.A. (2001), "Inducible expression of mutant α -synuclein decreases proteasome activity and increases sensitivity to mitochondria-dependent apoptosis", *Human Molecular Genetics*, Vol. 10, No. 9, pp. 919-926
- Thalmann, R., Henzl, M.T. and Thalmann, I. (1997), "Specific proteins of the organ of Corti", *Acta Otolaryngologica*, Vol. 117, No. 2, pp. 265-268

- Tsai, Y.C., Fishman, P.S., Thakor, N.V. and Oyler, G.A. (2003), "Parkin facilitates the elimination of expanded polyglutamine proteins and leads to preservation of proteasome function", *Journal of Biological Chemistry*, Vol. 278, No. 24, pp. 22044-22055
- Urushitani, M., Kurisu, J., Tateno, M., Hatakeyama, S., Nakayama, K., Kato, S. and Takahashi, R. (2004), "CHIP promotes proteasomal degradation of familial ALS-linked mutant SOD1 by ubiquitinating Hsp/Hsc70", *Journal of Neurochemistry*, Vol. 90, No. 1, pp. 231-244
- Varshavsky, A. (2000), "Recent studies of the Ubiquitin system and the N-end rule pathway", *Harvey lectures*, Vol. 96, No., pp. 93-116
- Vashist, S., Kim, W., Belden, W.J., Spear, E.D., Barlowe, C. and Ng, D.T.W. (2001), "Distinct retrieval and retention mechanisms are required for the quality control of endoplasmic reticulum protein folding", *Journal of Cell Biology*, Vol. 155, No. 4, pp. 355-367
- Verma, R., Chen, S., Feldman, R., Schieltz, D., Yates, J., Dohmen, J. and Deshaies, R.J. (2000), "Proteasomal proteomics: Identification of nucleotide-sensitive proteasome-interacting proteins by mass spectrometric analysis of affinity-purified proteasomes", *Molecular Biology of the Cell*, Vol. 11, No. 10, pp. 3425-3439
- Walker, W.P. and Gunn, T.M. (2010), "Piecing together the pigment-type switching puzzle", *Pigment Cell Melanoma Res*, Vol. 23, No. 1, pp. 4-6
- Wang, D., Qian, L., Xiong, H., Liu, J., Neckameyer, W.S., Oldham, S., Xia, K., Wang, J., Bodmer, R. and Zhang, Z. (2006), "Antioxidants protect PINK1-dependent dopaminergic neurons in *Drosophila*", *Proceedings of the National Academy of Sciences of the United States of America*, Vol. 103, No. 36, pp. 13520-13525
- Whatley, B.R., Li, L. and Chin, L.S. (2008), "The Ubiquitin-proteasome system in spongiform degenerative disorders", *Biochim Biophys Acta*, Vol. 1782, No. 12, pp. 700-712
- Wigley, C.W., Fabunmi, R.P., Lee, M.G., Marino, C.R., Muallem, S., DeMartino, G.N. and Thomas, P.J. (1999), "Dynamic association of proteasomal machinery with the centrosome", *Journal of Cell Biology*, Vol. 145, No. 3, pp. 481-490
- Wilhovsky, S., Gardner, R. and Hampton, R. (2000), "HRD gene dependence of endoplasmic reticulum-associated degradation", *Molecular Biology of the Cell*, Vol. 11, No. 5, pp. 1697-1708
- Winklhofer, K.F., Henn, I.H., Kay-Jackson, P.C., Heller, U. and Tatzelt, J. (2003), "Inactivation of parkin by oxidative stress and C-terminal truncations: A protective role of molecular chaperones", *Journal of Biological Chemistry*, Vol. 278, No. 47, pp. 47199-47208
- Wytenbach, A., Sauvageot, O., Carmichael, J., Diaz-Latoud, C., Arrigo, A.P. and Rubinsztein, D.C. (2002), "Heat shock protein 27 prevents cellular polyglutamine toxicity and suppresses the increase of reactive oxygen species caused by huntingtin", *Human Molecular Genetics*, Vol. 11, No. 9, pp. 1137-1151
- Xiong, H., Wang, D., Chen, L., Yeun, S.C., Ma, H., Tang, C., Xia, K., Jiang, W., Ronai, Z., Zhuang, X. and Zhang, Z. (2009), "Parkin, PINK1, and DJ-1 form a Ubiquitin E3 ligase complex promoting unfolded protein degradation", *Journal of Clinical Investigation*, Vol. 119, No. 3, pp. 650-660
- Xu, C., Bailly-Maitre, B. and Reed, J.C. (2005), "Endoplasmic reticulum stress: Cell life and death decisions", *Journal of Clinical Investigation*, Vol. 115, No. 10, pp. 2656-2664
- Yamanaka, K., Ishikawa, H., Megumi, Y., Tokunaga, F., Kanie, M., Rouault, T.A., Morishima, I., Minato, N., Ishimori, K. and Iwai, K. (2003), "Identification of the Ubiquitin-protein ligase that recognizes oxidized IRP2", *Nature Cell Biology*, Vol. 5, No. 4, pp. 336-340
- Yamanaka, T., Miyazaki, H., Oyama, F., Kurosawa, M., Washizu, C., Doi, H. and Nukina, N. (2008), "Mutant Huntingtin reduces HSP70 expression through the sequestration of NF-Y transcription factor", *EMBO Journal*, Vol. 27, No. 6, pp. 827-839
- Yang, H., Zhong, X., Ballar, P., Luo, S., Shen, Y., Rubinsztein, D.C., Monteiro, M.J. and Fang, S. (2007), "Ubiquitin ligase Hrd1 enhances the degradation and suppresses the toxicity of polyglutamine-expanded huntingtin", *Experimental Cell Research*, Vol. 313, No. 3, pp. 538-550
- Yashiro, K., Riday, T.T., Condon, K.H., Roberts, A.C., Bernardo, D.R., Prakash, R., Weinberg, R.J., Ehlers, M.D. and Philpot, B.D. (2009), "Ube3a is required for experience-dependent maturation of the neocortex", *Nature Neuroscience*, Vol. 12, No. 6, pp. 777-783
- Yasojima, K., Tsujimura, A., Mizuno, T., Shigeyoshi, Y., Inazawa, J., Kikuno, R., Kuma, K.I., Ohkubo, K., Hosokawa, Y., Ibata, Y., Abe, T., Miyata, T., Matsubara, K., Nakajima, K. and Hashimoto-Gotoh, T. (1997), "Cloning of human and mouse cDNAs encoding novel zinc finger proteins expressed in cerebellum and hippocampus", *Biochemical and Biophysical Research Communications*, Vol. 231, No. 2, pp. 481-487

- Ye, Y., Shibata, Y., Yun, C., Ron, D. and Rapoport, T.A. (2004), "A membrane protein complex mediates retro-translocation from the ER lumen into the cytosol", *Nature*, Vol. 429, No. 6994, pp. 841-847
- Yewdell, J.W., Antón, L.C. and Bennink, J.R. (1996), "Defective Ribosomal Products (DRiPs): A Major Source of Antigenic Peptides for MHC Class I Molecules?", *Journal of Immunology*, Vol. 157, No. 5, pp. 1823-1826
- Ying, Z., Wang, H., Fan, H., Zhu, X., Zhou, J., Fei, E. and Wang, G. (2009), "Gp78, an ER associated E3, promotes SOD1 and ataxin-3 degradation", *Human Molecular Genetics*, Vol. 18, No. 22, pp. 4268-4281
- Yoshida, Y., Adachi, E., Fukiya, K., Iwai, K. and Tanaka, K. (2005), "Glycoprotein-specific Ubiquitin ligases recognize N-glycans in unfolded substrates", *EMBO Reports*, Vol. 6, No. 3, pp. 239-244
- Yoshida, Y., Chiba, T., Tokunaga, F., Kawasaki, H., Iwal, K., Suzuki, T., Ito, Y., Matsuoka, K., Yoshida, M., Tanaka, K. and Tai, T. (2002), "E3 Ubiquitin ligase that recognizes sugar chains", *Nature*, Vol. 418, No. 6896, pp. 438-442
- Yoshida, Y., Tokunaga, F., Chiba, T., Iwai, K., Tanaka, K. and Tai, T. (2003), "Fbs2 Is a New Member of the E3 Ubiquitin Ligase Family That Recognizes Sugar Chains", *Journal of Biological Chemistry*, Vol. 278, No. 44, pp. 43877-43884
- Younger, J.M., Chen, L., Ren, H.Y., Rosser, M.F.N., Turnbull, E.L., Fan, C.Y., Patterson, C. and Cyr, D.M. (2006), "Sequential Quality-Control Checkpoints Triage Misfolded Cystic Fibrosis Transmembrane Conductance Regulator", *Cell*, Vol. 126, No. 3, pp. 571-582
- Zavacki, A.M., E Drigo, R.A., Freitas, B.C.G., Chung, M., Harney, J.W., Egri, P., Wittmann, G., Fekete, C., Gereben, B. and Bianco, A.C. (2009), "The E3 Ubiquitin ligase TEB4 mediates degradation of type 2 iodothyronine deiodinase", *Molecular and Cellular Biology*, Vol. 29, No. 19, pp. 5339-5347
- Zenker, M., Mayerle, J., Lerch, M.M., Tagariello, A., Zerres, K., Durie, P.R., Beier, M., Hälskamp, G., Guzman, C., Rehder, H., Beemer, F.A., Hamel, B., Vanlieferinghen, P., Gershoni-Baruch, R., Vieira, M.W., Dumic, M., Auslender, R., Gil-Da-Silva-Lopes, V.L., Steinlicht, S., Rauh, M., Shalev, S.A., Thiel, C., Winterpacht, A., Kwon, Y.T., Varshavsky, A. and Reis, A. (2005), "Deficiency of UBR1, a Ubiquitin ligase of the N-end rule pathway, causes pancreatic dysfunction, malformations and mental retardation (Johanson-Blizzard syndrome)", *Nature Genetics*, Vol. 37, No. 12, pp. 1345-1350
- Zhao, J., Ren, Y., Jiang, Q. and Feng, J. (2003), "Parkin is recruited to the centrosome in response to inhibition of proteasomes", *Journal of Cell Science*, Vol. 116, No. 19, pp. 4011-4019
- Zhong, X. and Pittman, R.N. (2006), "Ataxin-3 binds VCP/p97 and regulates retrotranslocation of ERAD substrates", *Human Molecular Genetics*, Vol. 15, No. 16, pp. 2409-2420
- Zhu, H., Kavsak, P., Abdollah, S., Wrana, J.L. and Thomsen, G.H. (1999), "A SMAD Ubiquitin ligase targets the BMP pathway and affects embryonic pattern formation", *Nature*, Vol. 400, No. 6745, pp. 687-693
- Zoghbi, H.Y. and Orr, H.T. (2000), "Glutamine repeats and neurodegeneration", *Annual Review of Neuroscience*, Vol. 23, No., pp. 217-247
- Zucchelli, S., Codrich, M., Marcuzzi, F., Pinto, M., Vilotti, S., Biagioli, M., Ferrer, I. and Gustincich, S. (2010), "TRAF6 promotes atypical ubiquitination of mutant DJ-1 and alpha-synuclein and is localized to Lewy bodies in sporadic Parkinson's disease brains", *Human Molecular Genetics*, Vol. 19, No. 19, pp. 3759-3770

...